## When Do Children Need Vaccinations?

Age	Hep B <sup>1</sup>	DTaP <sup>1</sup>	Polio <sup>1</sup>	MMR <sup>1</sup>	Varicella <sup>1+2</sup> (or health provider signed statement with dates of disease)	PCV <sup>2</sup>	Hib <sup>3</sup>	Rota Virus <sup>3</sup>	HepA <sup>3+4</sup>	Flu <sup>3</sup> (annually in fall for all over 6 months)
Birth	X									
1 mo	X									
2 mo	(1-2 mo)	X	X			X	X	X		
4 mo		X	X			X	X	X		
6 mo		X				X	X	X		×
12-18	×	×	X	X	X	X	X		X	
mo	(6-18 mo)	(15-18 mo)		(12-15 mo)		(12-15 mo)	(12-15 mo <b>)</b>		2 doses 6 months apart by age 24 mos	X
2-3										×
yr		~	<b>V</b>	~	<b>V</b>					~
4-6 yr		X	(4 <sup>th</sup> dose is recommended)	X	X					X

- 1 Required for Kindergarten and all child care types
- 2- Required for Centers, CCDF, VCP and Homes
- 3- Not required but highly recommended for children in child care
- 4- 2 doses required for Kindergarten entrance beginning 2014; highly recommended for child care

PCV 13 (pneumococcal conjugate, Prevnar)

	· congugues, · · · · · · · · · ·		
Recommended			
schedule for			
vaccination			
2 m			
4m			
6m			
12-15 m (booster)			
24-59m dose only if			
the child does not			
have 4 prior doses of			
PCV 13, or if the			
child completed 4			
doses of PCV 7			
(earlier version of			
vaccine).			