

Pressure Ulcer Prevention Points



NATIONAL
PRESSURE
ULCER
ADVISORY
PANEL

I. Risk Assessment

1. Consider all bed-bound and chair-bound persons, or those whose ability to reposition is impaired, to be at risk for pressure ulcers.
2. Use a valid, reliable and age appropriate method of risk assessment that ensures systematic evaluation of individual risk factors.
3. Assess all at-risk patients/residents at the time of admission to health care facilities, at regular intervals thereafter and with a change in condition. A schedule is helpful and should be based on individual acuity and the patient care setting.
 - Acute care: assess on admission, reassess at least every 24 hours or sooner if the patient's condition changes
 - Long-term care: assess on admission, weekly for four weeks, then quarterly and whenever the resident's condition changes
 - Home care: assess on admission and at every nurse visit.
4. Identify all individual risk factors (decreased mental status, exposure to moisture, incontinence, device related pressure, friction, shear, immobility, inactivity, nutritional deficits) to guide specific preventive treatments. Modify care according to the individual factors.
5. Document risk assessment subscale scores and total scores and implement a risk-based prevention plan.

II. Skin Care

1. Perform a head to toe skin assessment at least daily, especially checking pressure points such as sacrum, ischium, trochanters, heels, elbows, and the back of the head.
2. Individualize bathing frequency. Use a mild cleansing agent. Avoid hot water and excessive rubbing. Use lotion after bathing. For neonates and infants follow evidence-based institutional protocols
3. Establish a bowel and bladder program for patients

- with incontinence. When incontinence cannot be controlled, cleanse skin at time of soiling, and use a topical barrier to protect the skin. Select under pads or briefs that are absorbent and provide a quick drying surface to the skin. Consider a pouching system or collection device to contain stool and to protect the skin.
4. Use moisturizers for dry skin. Minimize environmental factors leading to dry skin such as low humidity and cold air. For neonates and infants follow evidence-based institutional protocols
 5. Avoid massage over bony prominences.

III. Nutrition

1. Identify and correct factors compromising protein/calorie intake consistent with overall goals of care.
2. Consider nutritional supplementation/support for nutritionally compromised persons consistent with overall goals of care.
3. If appropriate offer a glass of water when turning to keep patient/resident hydrated.
4. Multivitamins with minerals per physician's order.

IV. Mechanical Loading and Support Surfaces

1. Reposition bed-bound persons at least every two hours and chair-bound persons every hour consistent with overall goals of care.
2. Consider postural alignment, distribution of weight, balance and stability, and pressure redistribution when positioning persons in chairs or wheelchairs.
3. Teach chair-bound persons, who are able, to shift weight every 15 minutes.
4. Use a written repositioning schedule.
5. Place at-risk persons on pressure-redistributing mattress and chair cushion surfaces.
6. Avoid using donut-type devices and sheepskin for pressure redistribution.
7. Use pressure-redistributing devices in the operating room for individuals assessed to be at high risk for pressure ulcer development.

8. Use lifting devices (e.g., trapeze or bed linen) to move persons rather than drag them during transfers and position changes.
9. Use pillows or foam wedges to keep bony prominences, such as knees and ankles, from direct contact with each other. Pad skin subjected to device related pressure and inspect regularly.
10. Use devices that eliminate pressure on the heels. For short-term use with cooperative patients, place pillows under the calf to raise the heels off the bed. Place heel suspension boots for long-term use.
11. Avoid positioning directly on the trochanter when using the side-lying position; use the 30° lateral inclined position.
12. Maintain the head of the bed at or below 30° or at the lowest degree of elevation consistent with the patient's/resident's medical condition.
13. Institute a rehabilitation program to maintain or improve mobility/activity status.

V. Education

1. Implement pressure ulcer prevention educational programs that are structured, organized, comprehensive, and directed at all levels of health care providers, patients, family, and caregivers.
2. Include information on:
 - a. etiology of and risk factors for pressure ulcers
 - b. risk assessment tools and their application
 - c. skin assessment
 - d. selection and use of support surfaces
 - e. nutritional support
 - f. program for bowel and bladder management
 - g. development and implement individualized programs of skin care
 - h. demonstration of positioning to decrease risk of tissue breakdown
 - i. accurate documentation of pertinent data
3. Include mechanisms to evaluate program effectiveness in preventing pressure ulcers.

Pressure Ulcer Prevention Points



NATIONAL
PRESSURE
ULCER
ADVISORY
PANEL

REFERENCES

- Ayello EA. Preventing pressure ulcers and skin tears. In Mezey M, Fulmer T, Abraham I, Zwicker DA (editors). *Geriatric nursing protocols for best practice*. 2nd ed. New York: Springer Publishing Company, Inc. 2003. p 165-84.
- Ayello EA, Braden B. How and why to do pressure ulcer risk assessment. *Adv Skin Wound Care*. 2002 15(3): 125-131.
- Baharestani MM, Ratliff C and the National Pressure Ulcer Advisory Panel. Neonatal and pediatric pressure ulcers: an NPUAP white paper. *Adv Skin Wound Care*. 2007
- Black JM. Preventing heel pressure ulcers. *Nursing* 2004 Nov 34(11): 17
- Braden BJ, Makelbust J. Preventing pressure ulcers with the Braden scale: an update on this easy-to-use tool that assesses a patient's risk. *Am J Nurs* 105(6): 70-2.
- Brienza DM, Geyer MJ. Using support surfaces to manage tissue integrity. *Adv Skin Wound Care*. 2005 18(3): 151-7.
- Cullum N, Deeks J, Sheldon TA, Song F, Fletcher AW. *Cochrane Review. Beds, Mattresses and Cushions for Pressure Sore Prevention and Treatment*. In: The Cochrane Library, 2000. Issue 3.
- Posthauer, ME. Hydration: an essential nutrient. *Adv Skin Wound Care*. 2005 18(1): 32-3.
- Posthauer, ME, Jordan, RS, Sylvia, C. & National Pressure Ulcer Advisory Panel. Support Surface Initiative: Terms and Definitions and Patient Care. *Adv Skin Wound Care*, Nov 2006 19 (9): 487-9
- The National Pressure Ulcer Advisory Panel's Summary of the AHCPR Clinical Practice Guideline, Pressure Ulcers in Adults: Prediction and Prevention (AHCPR Publication No. 92-0047. Rockville, MD: May 1992). © 1993 NPUAP
- Wound Ostomy and Continence Nurses Society. (2003) *Guideline for Prevention and Management of Pressure Ulcers*. Glenview: IL. WOCN.

ADDITIONAL REFERENCES

- Ankrom MA, Bennett RG, Sprigle S, Langemo D, Black JM, Berlowitz DR, Lyder CH. National Pressure Ulcer Advisory Panel. Pressure-related deep tissue injury under intact skin and the current pressure ulcer staging systems. *Adv Skin Wound Care*. 2005 Jan-Feb 18 (1): 35-42
- Berlowitz DR, Ratliff C, Cuddigan J, Rodeheaver GT; National Pressure Ulcer Advisory Panel. The PUSH Tool: A Survey to Determine Its Perceived Usefulness. *Adv Skin Wound Care*. 2005 Nov-Dec; 18(9): 480-483.
- Black JM and the National Pressure Ulcer Advisory Panel. Moving Toward Consensus on Deep Tissue Injury and Pressure Ulcer Staging. *Adv Skin Wound Care*. 2005 Oct; 18(8): 415-421.
- Cuddigan J, Frantz RA. Pressure ulcer research: pressure ulcer treatment. A monograph from the National Pressure Ulcer Advisory Panel. *Adv Skin Wound Care*. 1998 Oct 11(6): 294-300; quiz 30.
- Henderson CT, Ayello EA, Sussman C, Leiby DM, Bennett MA, Dungog EF, Sprigle S, Woodruff L. Draft definition of stage I pressure ulcers: inclusion of persons with darkly pigmented skin. NPUAP Task Force on Stage I Definition and Darkly Pigmented Skin. *Adv Skin Wound Care*. 1997 Sept; 10(5): 16-9.
- National Pressure Ulcer Advisory Panel. Pressure ulcers in America: prevalence, incidence, and implications for the future. An executive summary of the National Pressure Ulcer Advisory Panel Monograph. *Adv Skin Wound Care*. 2001 Jul-Aug. 14(4): 208-15.
- National Pressure Ulcer Advisory Panel. *Monograph Pressure Ulcers in America: Prevalence, Incidence, and Implications for the Future*. April 1, 2001.
- National Pressure Ulcer Advisory Panel. Management of tissue load: An excerpt from the third NPUAP Slide Set. National Pressure Ulcer Advisory Panel. *Adv Skin Wound Care*. 1997 Oct. 10(6): 35-8.
- National Pressure Ulcer Advisory Panel. Monitoring Pressure Ulcer Healing: An Alternative to Reverse Staging. Proceedings of the National Pressure Ulcer Advisory Panel 5th National Conference. 1997. *Adv Skin Wound Care*. 1997 Sept. 10(5): 8-10.
- National Pressure Ulcer Advisory Panel. Description of NPUAP. National Pressure Ulcer Advisory Panel. *Adv Skin Wound Care*. Jul-Aug 8(4): suppl 93-5.
- National Pressure Ulcer Advisory Panel. Pressure ulcer prevalence, cost and risk assessment: consensus development conference statement—The National Pressure Ulcer Advisory Panel. *Decubitus* 1989 May 2(2): 24-8.
- Pieper B. Mechanical forces: pressure, shear and friction. In *Acute and Chronic Wounds*, R, Bryant and D Nix. Ed. 2007: 205-234.
- Thomas DR, Rodeheaver GT, Bartolucci AA, Franz RA, Sussman C, Ferrell BA, Cuddigan J, Stotts NA, Makelbust J. Pressure ulcer scale for healing: derivation and validation of the PUSH tool. The PUSH Task Force. *Adv Skin Wound Care*. 1997 10(5): 96-101.
- Zanca JM, Brienza DM, Berlowitz D, Bennett RG, Lyder CH; National Pressure Ulcer Advisory Panel. Pressure ulcer research funding in America: creation and analysis of an on-line database. *Adv Skin Wound Care*. 2003 Jul-Aug; 16(4): 190-7.
- Zulkowski K, Langemo D, Posthauer ME; National Pressure Ulcer Advisory Panel. Coming to consensus on deep tissue injury. *Adv Skin Wound Care*. 2005 Jan-Feb; 18(1): 28-9.

National Pressure Ulcer Advisory Panel
1255 Twenty-Third Street NW, Suite 200
Washington, DC 20037

T: 202-521-6789
F: 202-833-3636
www.npuap.org