

STD MORBIDITY

Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2018

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	23,473	68.1%	5,992	49.9%	23	6.3%
Male	10,990	31.9%	6,019	50.1%	341	93.7%
Unknown	18	<1%	9	<1%	0	0%
Total	34,481	100%	12,020	100%	364	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	14,572	42.3%	4,767	40.0%	175	48.1%
Black	11,559	33.5%	5,396	44.9%	152	41.8%
All Others	3,989	11.5%	918	7.3%	35	10.1%
Unknown	4,361	12.7%	939	7.8%	2	<1%
Total	34,481	100%	12,020	100%	364	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	2,576	7.5%	503	4.2%	26	7.1%
Non-Hispanic	26,703	77.5%	10,264	85.4%	338	92.9%
Unknown	5,202	15.1%	1,253	10.4%	0	<1%
Total	34,481	100%	12,020	100%	364	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	10,231	29.7%	2,255	18.8%	16	4.4%
20-24	12,970	37.6%	3,521	29.3%	80	22.0%
25-29	6,100	17.7%	2,744	22.8%	85	23.4%
30-34	2,606	7.6%	1,519	12.6%	53	14.6%
35-39	1,371	4.0%	917	7.6%	45	12.4%
40-44	613	1.8%	434	3.6%	20	5.4%
45-54	459	1.3%	468	3.9%	43	11.8%
55+	131	<1%	162	1.4%	22	6.0%
Unknown	0	0%	0	0%	0	0%
Total	34,481	100%	12,020	100%	364	100%

Source: ISDH STD Database as of February 7, 2019

NOTE: Not all percents may add to 100 due to rounding.