

STD MORBIDITY

Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2013

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	20,307	72.4%	3,796	53.1%	18	8.4%
Male	7,708	27.5%	3,347	46.8%	197	91.6%
Unknown	8	<1%	1	<1%	0	0
Total	28,023	100%	7,144	100%	215	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	13,096	46.7%	2,354	33.0%	93	43.3%
Black	10,191	36.4%	3,988	55.8%	89	41.4%
All Others	4,736	16.9%	802	11.2%	33	15.3%
Unknown	0	0	0	0	0	0
Total	28,023	100%	7,144	100%	215	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	1,564	5.6%	233	3.3%	23	10.7%
Non-Hispanic	26,459	94.4%	6,342	88.8%	182	84.6%
Unknown	0	0	569	7.9%	10	4.7%
Total	28,023	100%	7,144	100%	215	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	8,851	31.6%	1,774	24.8%	14	6.5%
20-24	11,365	40.5%	2,561	35.8%	64	29.8%
25-29	4,382	15.6%	1,345	18.8%	41	19.1%
30-34	1,884	6.7%	650	9.1%	31	14.4%
35-39	819	2.9%	341	4.8%	19	8.8%
40-44	374	1.3%	206	2.9%	11	5.1%
45-54	279	1%	209	2.9%	28	13.0%
55+	69	<1%	58	<1%	7	3.3%
Unknown	0	0	0	0	0	0
Total	28,023	100%	7,144	100%	215	100%

Source: ISDH STD Database as of January 22, 2015

NOTE: Not all percents may add to 100 due to rounding.