

# STD MORBIDITY

## Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2012

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	21,633	73.3%	4,139	56.4%	22	9.8%
Male	7,850	26.6%	3,188	43.4%	202	90.2%
Unknown	22	<1%	11	<1%		
Total	29,505	100%	7,338	100%	224	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	12,566	42.6%	2,080	28.3%	98	43.8%
Black	10,410	35.3%	4,087	55.7%	110	49.1%
Amer. Indian	41	<1%	12	<1%		
Asian	86	<1%	11	<1%	1	<1%
Multiple Races	251	<1%	63	<1%	3	1.3%
All Others	6,151	20.8%	1,085	14.8%	12	5.4%
Unknown						
Total	29,505	100%	7,338	100%	224	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	1,371	4.6%	208	2.8%	7	3.1%
Non-Hispanic	28,134	95.4%	7,130	97.2%	217	96.9%
Unknown						
Total	29,505	100%	7,338	100%	224	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	10,089	34.2%	1,885	25.7%	32	14.3%
20-24	11,579	39.2%	2,671	36.4%	53	23.7%
25-29	4,355	14.8%	1,330	18.1%	44	19.6%
30-34	1,902	6.4%	675	9.2%	21	9.4%
35-39	773	2.6%	338	4.6%	16	7.1%
40-44	404	1.4%	204	2.8%	26	11.6%
45-54	310	1%	179	2.4%	26	11.6%
55+	93	<1%	56	<1%	6	2.7%
Unknown						
Total	29,505	100%	7,338	100%	224	100%

Source: ISDH STD Database as of August 8, 2013

NOTE: Not all percents may add to 100 due to rounding.