


# 1st Congressional District Nutrient and Sediment Load Reductions


## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District




Land use calculated using the 2020 NASS Cropland Data Layer



- County Seats
- County Boundaries
- Congressional District

### Sediment Reduced: 31,635,165 lbs.

Enough to fill 158 freight cars!



### Phosphorus Reduced: 20,170 lbs.

Enough to fill 20 truck beds (8' bed)!



### Nitrogen Reduced: 40,285 lbs.

Enough to fill 40 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	153	219	16,467,295	10,085	20,180
2016	110	183	12,406,370	7,210	14,380
2017	131	230	18,701,100	11,625	23,200
2018	155	274	18,629,325	11,585	23,110
2019	340	494	32,633,600	20,740	41,420
2020	256	437	31,635,165	20,170	40,285
<b>13-20</b>	<b>1,441</b>		<b>161,014,425</b>	<b>100,755</b>	<b>201,235</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)  
Last updated: 4/7/2021


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

# 2nd Congressional District Nutrient and Sediment Load Reductions


## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District




Land use calculated using the 2020 NASS Cropland Data Layer



### Sediment Reduced: 204,473,285 lbs.

Enough to fill 1,022 freight cars!



### Phosphorus Reduced: 111,155 lbs.

Enough to fill 111 truck beds (8' bed)!



### Nitrogen Reduced: 222,425 lbs.

Enough to fill 222 truck beds (8' bed)!




Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	1,708	2,081	216,213,330	123,625	248,270
2016	1,435	1,913	188,357,040	106,110	212,520
2017	1,581	2,286	198,693,845	111,930	224,225
2018	1,366	2,211	168,465,780	95,495	191,265
2019	1,480	2,558	211,005,765	113,575	227,220
2020	1,994	3,340	204,473,285	111,155	222,425
<b>13-20</b>	<b>13,324</b>		<b>1,641,622,980</b>	<b>920,235</b>	<b>1,843,275</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

# 3rd Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District



Land use calculated using the 2020 NASS Cropland Data Layer

### Sediment Reduced: 364,878,565 lbs.

Enough to fill 1,824 freight cars!



### Phosphorus Reduced: 234,945 lbs.

Enough to fill 235 truck beds (8' bed)!



### Nitrogen Reduced: 501,195 lbs.

Enough to fill 501 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	2,448	3,031	309,059,770	190,600	395,075
2016	2,055	2,886	287,297,795	180,195	374,260
2017	2,526	3,506	322,691,775	200,500	414,845
2018	2,712	4,023	290,218,305	192,035	415,490
2019	3,615	5,190	443,066,670	281,570	594,395
2020	2,929	4,861	364,878,565	234,945	501,195
<b>13-20</b>	<b>21,295</b>		<b>2,603,550,020</b>	<b>1,628,755</b>	<b>3,420,965</b>

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)  
 Last updated: 4/7/2021


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

# 4th Congressional District Nutrient and Sediment Load Reductions


## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District




Land use calculated using the 2020 NASS Cropland Data Layer



### Sediment Reduced: 488,867,810 lbs.

Enough to fill 2,444 freight cars!



### Phosphorus Reduced: 275,910 lbs.

Enough to fill 276 truck beds (8' bed)!



### Nitrogen Reduced: 558,680 lbs.

Enough to fill 559 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	1,751	3,020	295,670,170	171,210	344,910
2016	1,533	3,266	272,139,170	156,935	322,265
2017	1,727	3,952	331,412,525	191,195	390,140
2018	2,002	4,495	372,281,570	213,170	433,680
2019	1,870	4,784	394,496,535	223,930	454,975
2020	2,641	5,917	488,867,810	275,910	558,680
<b>13-20</b>	<b>14,873</b>		<b>2,571,875,055</b>	<b>1,478,880</b>	<b>3,000,865</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

# 5th Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership




### Comparison of Landuse Across District



Land use calculated using the 2020 NASS Cropland Data Layer

### Sediment Reduced: 39,986,525 lbs.

Enough to fill 200 freight cars!



### Phosphorus Reduced: 26,420 lbs.

Enough to fill 26 truck beds (8' bed)!



### Nitrogen Reduced: 53,830 lbs.

Enough to fill 54 truck beds (8' bed)!




Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	189	268	27,069,660	18,160	37,350
2016	73	225	15,331,050	9,790	20,615
2017	142	315	23,450,775	14,770	30,565
2018	162	378	24,146,555	15,320	31,670
2019	206	444	34,873,295	22,490	45,995
2020	203	465	39,986,525	26,420	53,830
<b>13-20</b>	<b>1,392</b>		<b>209,989,560</b>	<b>136,160</b>	<b>278,395</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

# 6th Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership




### Comparison of Landuse Across District



Land use calculated using the 2020 NASS Cropland Data Layer

### Sediment Reduced: 553,350,595 lbs.

Enough to fill 2,767 freight cars!



### Phosphorus Reduced: 283,095 lbs.

Enough to fill 283 truck beds (8' bed)!



### Nitrogen Reduced: 567,085 lbs.

Enough to fill 567 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	1,511	2,338	295,058,725	148,135	296,720
2016	1,554	2,742	297,080,235	149,175	298,660
2017	1,476	3,097	301,723,640	153,530	307,360
2018	2,026	3,943	371,381,825	191,380	383,700
2019	2,656	5,152	483,447,925	250,025	500,820
2020	2,802	5,678	553,350,595	283,095	567,085
<b>13-20</b>	<b>14,984</b>		<b>2,777,670,300</b>	<b>1,414,685</b>	<b>2,833,525</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

# 7th Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District



Land use calculated using the 2020 NASS Cropland Data Layer



### Sediment Reduced: 1,668,880 lbs.

Enough to fill 8 freight cars!



### Phosphorus Reduced: 695 lbs.

Enough to fill 0.70 truck beds (8' bed)!



### Nitrogen Reduced: 1,335 lbs.

Enough to fill 1.34 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	20	31	2,189,850	1,030	1,995
2016	33	45	2,018,970	920	1,770
2017	36	58	2,176,835	1,040	2,015
2018	29	57	1,717,210	725	1,380
2019	23	60	1,830,925	790	1,525
2020	25	77	1,668,880	695	1,335
<b>13-20</b>	<b>199</b>		<b>14,137,170</b>	<b>6,445</b>	<b>12,445</b>


The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)  
 Last updated: 4/7/2021


Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

# 8th Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District




Land use calculated using the 2020 NASS Cropland Data Layer



### Sediment Reduced: 1,022,563,42 lbs.

Enough to fill 5,113 freight cars!



### Phosphorus Reduced: 482,570 lbs.

Enough to fill 483 truck beds (8' bed)!



### Nitrogen Reduced: 965,875 lbs.

Enough to fill 966 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.


Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	2,730	4,047	856,394,375	400,300	801,075
2016	2,254	3,987	746,895,275	350,235	701,180
2017	2,426	4,697	830,417,050	388,760	778,485
2018	3,638	6,226	901,513,620	422,710	846,365
2019	3,308	6,887	957,661,750	455,185	911,275
2020	3,792	7,943	1,022,563,425	482,570	965,875
<b>13-20</b>	<b>23,880</b>		<b>6,787,408,660</b>	<b>3,188,955</b>	<b>6,383,525</b>

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.




# 9th Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



### Comparison of Landuse Across District



Land use calculated using the 2020 NASS Cropland Data Layer

**Sediment Reduced: 657,090,885 lbs.**

Enough to fill 3,285 freight cars!



**Phosphorus Reduced: 307,855 lbs.**

Enough to fill 308 truck beds (8' bed)!



**Nitrogen Reduced: 618,590 lbs.**

Enough to fill 619 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2015	1,500	2,295	534,566,110	250,165	502,825
2016	1,650	2,840	568,840,840	264,470	531,470
2017	1,865	3,462	703,934,560	326,905	656,425
2018	2,004	3,955	632,092,045	295,120	593,185
2019	2,392	4,765	666,883,830	311,965	626,905
2020	2,508	5,305	657,090,885	307,855	618,590
<b>13-20</b>	<b>14,751</b>		<b>4,496,782,515</b>	<b>2,094,775</b>	<b>4,206,180</b>

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5. Please Note: Calendar year 2013 and 2014 metrics are excluded from the table due to space limitations, but are present in the "13-20" summations.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)  
Last updated: 4/7/2021

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.