





9th Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Sediment Reduced: 632,488,395 lbs.
Enough to fill 3,162 freight cars!



Phosphorus Reduced: 295,325 lbs.
Enough to fill 295 truck beds (8' bed)!



Nitrogen Reduced: 593,590 lbs.
Enough to fill 593 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	1,492	1,492	322,690,665	148,920	297,940
2014	1,340	1,835	410,683,580	189,375	378,840
2015	1,500	2,295	534,566,110	250,165	502,825
2016	1,650	2,850	569,237,190	264,675	531,875
2017	1,865	3,472	704,330,910	327,110	656,835
2018	2,004	3,965	632,488,395	295,325	593,590
13-18	9,851		3,173,996,850	1,475,570	2,961,905

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5.