



# Geist Reservoir Watershed Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



**Sediment Reduced: 1,414,400 lbs.**

Enough to fill 7 freight cars!



**Phosphorus Reduced: 925 lbs.**

Enough to fill 1 truck beds (8' bed)!



**Nitrogen Reduced: 1,895 lbs.**

Enough to fill 2 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	16	16	2,596,730	1,720	3,435
2014	29	29	2,750,285	1,855	3,705
2015	11	25	972,180	615	1,270
2016	6	23	511,690	345	725
2017	18	35	1,779,975	1,185	2,415
2018	14	34	1,414,400	925	1,895
<b>13-18</b>	<b>94</b>		<b>10,025,260</b>	<b>6,645</b>	<b>13,445</b>

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5.

For more information visit: <http://www.in.gov/isda/2991.htm> or contact [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)  
 Last updated: 3/8/2019

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.