

STD MORBIDITY

Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2014

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	20,586	72.2%	3,819	52.4%	11	6.5%
Male	7,921	27.8%	3,465	47.5%	157	93.5%
Unknown	12	<1%	5	<1%	0	0%
Total	28,519	100%	7,289	100%	168	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	13,851	48.6%	2,577	35.3%	75	44.6%
Black	10,633	37.3%	4,109	56.4%	79	47.0%
All Others	4,035	14.1%	603	8.3%	14	8.4%
Unknown	0	0%	0	0%	0	0%
Total	28,519	100%	7,289	100%	168	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	1,693	5.9%	220	3.0%	8	4.8%
Non-Hispanic	26,826	94.1%	7069	97.0%	160	95.2%
Unknown	0	0%	0	0%	0	0%
Total	28,519	100%	7,289	100%	168	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	8,426	29.5%	1,664	22.8%	4	2.4%
20-24	11,698	41.0%	2,651	36.4%	52	30.9%
25-29	4,704	16.5%	1,359	18.6%	35	20.8%
30-34	1,979	6.9%	722	9.9%	18	10.7%
35-39	898	3.1%	395	5.4%	21	12.5%
40-44	424	1.5%	208	2.8%	10	6.0%
45-54	316	1.1%	214	2.9%	24	14.3%
55+	73	<1%	75	1%	4	2.4%
Unknown	1	<1%	1	<1%	0	0%
Total	28,519	100%	7,289	100%	168	100%

Source: ISDH STD Database as of November 4, 2015

NOTE: Not all percents may add to 100 due to rounding.