

Care Coordination Services

Currently there are 18 Care Coordination site in Indiana that provide health and human services for people living with HIV/AIDS (2005).

This report will present the demographic characteristics of the persons that use those services. In Table 65, the users of care services are broken out by sex and the quarter in which they used some of Care Coordination services.


Table 67: Care Utilization Services by Gender and Quarter in Percent, July 2005 – June 2006.

	July- September 2005	October- December 2005	January – March 2006	April – June 2006	Average
Male	77.9%	77.6%	78.1%	78.2%	77.9%
Female	22.1%	22.4%	21.9%	21.8%	22.1%
Total	3,137	3,249	3,357	3,280	3,256

Consistent with the HIV infected population at large, more than three-quarters of recipients were male in 2004.

In terms of race and ethnicity, the distribution of care recipients is shown in Figure 74.

Figure 74: Care Utilization Services by Race/Ethnicity and Quarter, 2005



The majority of service recipients were White (57.7%), followed by Black/African-Americans (30.8%). The share of Hispanics (7.6%) among Care Utilization users is almost twice their share


of the general population in Indiana. Table 66 below lists the racial and ethnic percentages by quarter and by race/ethnicity.

Table 68: Care Utilization Services by Race/Ethnicity and Quarter, 2005

Race	July-September 2005	October-December 2005	January – March 2006	April-June 2006	Average
Black	30.6%	30.3%	31.4%	30.9%	30.8%
White	59.6%	59.4%	53.1%	58.8%	57.7%
Hispanic	6.0%	6.2%	11.5%	6.7%	7.6%
Other	3.8%	4.2%	4.0%	3.6%	3.9%
Total	3,137	3,249	3,357	3,280	3,256

Figure 75 shows the percentages of Care Coordination services recipients by age group for 2005.


Figure 75: Percentage of Care Coordinator Clients by Age Category, 2005



The majority of the 3,280 clients receiving care services in 2005 were in the age range of 40 to 49 years of age.

The distribution of risk categories among the service recipients is shown in Figure 76.

Figure 76: Care Utilization Services by Exposure and Quarter for 2005



More than half of all recipients are associated with MSM as their primary risk category, consistent with the risk category distribution of the HIV/AIDS infected population overall.

Table 67 lists the corresponding percentages and totals by risk category and quarter for 2005.

Table 69: Care Utilization Services by Exposure and Quarter for 2005

Exposure	July - September 2005	October- December 2005	January – March 2006	April- June 2006	Average
MSM	52.5%	51.5%	52.5%	53.5%	52.5%
IDU	5.7%	5.9%	5.7%	5.6%	5.7%
MSM/IDU	2.4%	2.3%	4.3%	2.3%	2.8%
Hetero	27.5%	27.5%	26.2%	28.2%	27.4%
Mother	1.2%	1.3%	1.4%	1.4%	1.3%
Other	10.7%	11.5%	9.9%	9.0%	10.3%
Total	3,137	3,249	3,357	3,280	3,356