

Question 2

What is the scope of the HIV/AIDS Epidemic in Indiana?

Overall HIV/AIDS Trends

Throughout this report, unless mentioned otherwise, the time period that is reported on covers the 12-month period from January 1, 2007 to December 31, 2007.

At the end of December 2007, Indiana had a total of 4,025 residents diagnosed with HIV, and 4,826 residents that had developed AIDS. That adds up to a total 8,851 persons living with HIV/AIDS in Indiana by the end of 2007. This number, also called the prevalence number, is the number of all persons that have been diagnosed with the HIV virus sometime in the past, are either HIV positive only, or have developed AIDS, and are still living at the cut-off time for data collection for this report. In comparison, the number of HIV/AIDS diagnosed people at the end of 2006 was 7,555 persons. That is the equivalent of a 14.6% increase in the number of HIV/AIDS diagnosed persons over the past year.

In order to be able to compare the absolute numbers of cases with other entities such as surrounding states or the nation as a whole, the absolute numbers of diagnosed persons will be converted into rates, in this case a prevalence rate per 100,000 persons of the specific population. By dividing the number of diagnosed persons by the total number of the population the HIV prevalence rate per 100,000 comes to 63.8 (58.0), for AIDS it calculates to 76.4 (63.9), and for HIV/AIDS combined it is 140.2 (121.9) per 100,000 persons. The numbers in parenthesis show the diagnosis rates at the end of 2006. The overall increase of 14.6% from the previous year is also reflected in the individual rate increases for 2007.

Within the United States, Indiana ranked 26th by number of cumulative AIDS cases reported in 2006, the last year for which state comparison data were available. Compared to its neighboring states, Table 6 shows a pulled-out section of the larger Table 5 on the cumulative number of reported AIDS cases for Indiana.

Table 5: Cumulative Number of AIDS Cases by State, Reported Through December 2006³

Rank	State	AIDS Cases	Rank	State	AIDS Cases
	United States	992,865	28	Oregon	6,015
1	New York	177,262	29	Nevada	5,762
2	California	142,918	30	Oklahoma	4,862
3	Florida	105,614	31	Minnesota	4,845
4	Texas	70,127	32	Kentucky	4,632
5	New Jersey	49,528	33	Wisconsin	4,546
6	Illinois	33,902	34	Arkansas	3,927
7	Pennsylvania	33,782	35	Delaware	3,573
8	Georgia	31,965	36	Hawaii	2,944
9	Maryland	30,571	37	Kansas	2,795
10	Puerto Rico	29,911	38	New Mexico	2,610
11	Massachusetts	19,395	39	Rhode Island	2,608
12	Louisiana	17,740	40	Utah	2,315
13	District of Columbia	17,561	41	Iowa	1,740
14	Virginia	16,979	42	West Virginia	1,511
15	North Carolina	16,072	43	Nebraska	1,490
16	Ohio	15,095	44	Maine	1,118
17	Michigan	15,054	45	New Hampshire	1,084
18	Connecticut	14,899	46	Alaska	655
19	South Carolina	13,406	47	Virgin Islands	647
20	Tennessee	12,516	48	Idaho	606
21	Washington	11,826	49	Vermont	467
22	Missouri	11,077	50	Montana	376
23	Arizona	10,442	51	South Dakota	260
24	Colorado	8,773	52	Wyoming	230
25	Alabama	8,702	53	North Dakota	145
26	Indiana	8,295	54	Guam	69
27	Mississippi	6,698			

³ Kaiser Family Foundation, *Cumulative Number of AIDS Cases, Reported Through December 2006*

Table 6: Cumulative Number of AIDS Cases, Reported Through December 2006, Selected Midwestern States and the U.S.⁴

Rank	State	AIDS Cases	Rate/100,000*
32	Kentucky	4,632	113.77
18	Ohio	15,095	133.51
26	Indiana	8,295	131.38
17	Michigan	15,054	151.00
6	Illinois	33,902	269.00
	United States	992,865	335.36

In Table 6 the states were arranged in ascending order according to the rate/100,000 persons of their respective population. Rank refers to the ranking by absolute number of diagnosed people among all states in the nation. There are differences between the ranking by absolute numbers and by rates as shown. For example, Indiana has a similar rate (131.38/100,000) as Ohio (133.51/100,000), even though Ohio has more cases in absolute numbers. Compared to the nation as a whole, Indiana ranks in midfield by absolute numbers (26th in the nation).


⁴ Kaiser Family Foundation, *Cumulative Number of HIV/AIDS Cases, Reported Through December 2004*

Prevalence (living only) of HIV/AIDS in Indiana

Prevalence numbers describe the number of cases of a disease in a population up to a certain point in time. In the case of this report, prevalence describes the number of persons diagnosed with HIV/AIDS in Indiana that were alive by December 31, 2007 and that were reported in the HIV/AIDS Surveillance Report.

The prevalence rate for HIV/AIDS in Indiana shows some significant details when breaking out the rate by gender, race/ethnicity or age. Figure 13 shows the HIV/AIDS rates by sex.

Figure 13: Prevalence Rates for HIV, AIDS, and HIV/AIDS by Sex, Indiana 2007



The rate of diagnosed males per 100,000 people of the overall male population is at 218.2 persons per 100,000, up from 200.5 in 2006. Males have an almost 4.5 times larger HIV/AIDS prevalence rate than females do. The female prevalence rate is at 51.1 persons per 100,000 females of the overall population, also up from 45.9 in 2006. For HIV and AIDS separately, males continue to be affected much more strongly than their female counterparts. HIV/AIDS continues to affect more males than females. Table 7 breaks out the absolute numbers, percentages and rates by sex and disease status.


Table 7: Prevalence Numbers, Percentages and Rates for HIV, AIDS, and HIV/AIDS by Sex, 2007

	HIV			AIDS			HIV/AIDS		
	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000
Male	3123	77.6	100.4	4022	83.3	129.3	7145	80.7	229.7
Female	902	22.4	28.2	804	16.7	25.1	1706	19.3	53.3
Total	4025	100.0	63.8	4826	100.0	76.4	8851	100.0	140.2


Prevalence of HIV/AIDS by Age

In order to better understand the dynamics at play with diagnosed persons it is helpful to look at two different age definitions. One is the age of diagnosed persons at the end of December 2007. The other is the age distribution for persons at the time of their diagnosis. Figure 14 shows the age group distribution for diagnosed persons that were living at the end of 2007.

Figure 14: Prevalence Rates for HIV/AIDS by Current Age in December 2007




The majority of diagnosed persons are in the groups of 30 to 39 years old and 40 to 49 years of age. In comparison, Figure 15 shows the age group distribution by age at time of diagnosis.

Figure 15: Prevalence Rates for HIV/AIDS by Age at Time of Diagnosis, 2007

It is interesting to note the difference in the age distribution between the two age profiles. Whereas the majority of diagnosed persons living at the end of 2007 were in their thirties and forties, a majority of persons were diagnosed in their twenties and thirties. For one, this discrepancy in the age distribution shows the effects of the availability and effectiveness of HIV/AIDS medications that have increased the long-term survival rate of diagnosed persons. In other words, diagnosed persons are living longer because of more effective medications, which is reflected in the greater number of diagnosed persons in the older age groups. As a group, the persons that have been diagnosed longer ago have moved from the age group of their initial diagnosis to their current age group at the end of 2007. In contrast, the majority of new diagnoses still occur in the age groups of 25 to 29 years of age and 30 to 39 years of age. As the group of the “initial” diagnoses is aging, the differences in age at time of diagnosis and age at the time of this study continues to grow.

Similar to this finding is the breakout by HIV and AIDS by age group. Figure 16 shows the corresponding distributions.


Figure 16: Prevalence Rates for HIV and AIDS by Current Age, 2007



The majority of diagnosed persons are currently living with HIV in their early thirties and forties. The majority of AIDS diagnosis are currently living in their late thirties and forties. The figure above gives a snapshot of the age distribution of the diagnosed population at the end of 2007.

The corresponding diagnoses ages for HIV and AIDS are in Figure 17. They show the majority of persons diagnosed with HIV are in their early twenties and thirties. The majority of persons diagnosed with AIDS are in their late twenties and thirties.

Figure 17: Prevalence Rates for HIV and AIDS by Age at Time of Diagnosis, 2007



The absolute numbers, percentages and rates for HIV, AIDS and HIV/AIDS by age at end of study are presented in Table 8 and for age at time of diagnosis in Table 9.

Table 8: Prevalence Numbers, Percentages and Rates for HIV, AIDS, and HIV/AIDS by Current Age, 2007

Age	HIV			AIDS			HIV/AIDS		
	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000
0 to 9	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
10 to 14	25	0.6	5.7	16	0.3	3.6	41	0.5	9.3
15 to 19	26	0.6	5.8	13	0.3	2.9	39	0.4	8.7
20 to 24	192	4.8	43.5	50	1.0	11.3	242	2.7	54.8
25 to 29	379	9.4	85.4	168	3.5	37.9	547	6.2	123.3
30 to 39	1118	27.8	133.8	1051	21.8	125.8	2169	24.5	259.7
40 to 49	1480	36.8	156.1	2220	46.0	234.1	3700	41.8	390.2
50 to 59	608	15.1	73.4	1038	21.5	125.4	1646	18.6	198.8
60 to 64	111	2.8	39.5	167	3.5	59.5	278	3.1	99.0
over 65	86	2.1	11.0	103	2.1	13.1	189	2.1	24.1
Total	4025	100.0	63.8	4826	100.0	76.4	8851	100.0	140.2

Note that 10-19 and 20-29 are split into two age groups.

Table 9: Prevalence Numbers, Percentages and Rates for HIV, AIDS, and HIV/AIDS by Age at Time of Diagnosis, 2007

Age	HIV			AIDS			HIV/AIDS		
	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000
0 to 9	16	0.4	1.9	9	0.2	1.0	25	0.3	2.9
10 to 14	32	0.8	7.2	23	0.5	5.2	55	0.6	12.4
15 to 19	195	4.8	43.3	43	0.9	9.5	238	2.7	52.8
20 to 24	655	16.3	148.3	256	5.3	58.0	911	10.3	206.3
25 to 29	794	19.7	179.0	690	14.3	155.5	1484	16.8	334.5
30 to 39	1445	35.9	173.0	2156	44.7	258.1	3601	40.7	431.1
40 to 49	658	16.3	69.4	1233	25.5	130.0	1891	21.4	199.4
50 to 59	176	4.4	21.3	353	7.3	42.6	529	6.0	63.9
60 to 64	25	0.6	8.9	44	0.9	15.7	69	0.8	24.6
over 65	29	0.7	3.7	19	0.4	2.4	48	0.5	6.1
Total	4025	100.0	63.8	4826	100.0	76.4	8851	100.0	140.2


Note that 10-19 and 20-29 are split into two age groups.

Prevalence of HIV/AIDS by Race/Ethnicity

A look at the racial and ethnic make-up provides further details on the composition of the diagnosed population.


Indiana is overwhelmingly white and Non-Hispanic. Figure 18 shows the prevalence rate by race/ethnicity at the end of 2007. The racial and ethnic composition of diagnosed persons in Indiana differs to a great extent from the racial/ethnic distribution of the state's overall population.

Figure 18: Prevalence Rates of HIV/AIDS by Race/Ethnicity in 2007



The overwhelming majority of diagnosed cases are among Blacks, even though Blacks account for only about 8.9% of the overall population. In other words, HIV/AIDS is overwhelming more prevalent among Blacks than any other racial or ethnic group. The separate view of HIV and AIDS reveals further details about racial and ethnic differences among the diagnosed population.

Figure 19: Prevalence Rates of HIV and AIDS by Race/Ethnicity in 2007



All of the racial groups have slightly higher rates of AIDS diagnosis than HIV diagnosis. Table 10 lists the numbers, percentages and rates for HIV, AIDS and HIV/AIDS by race/ethnicity.


Table 10: Prevalence Numbers, Percentages and Rates of HIV, AIDS, and HIV/AIDS by Race/Ethnicity in 2007

Race/ Ethnicity	HIV			AIDS			HIV/AIDS		
	Number	%	Rate	Number	%	Rate	Number	%	Rate
Black	1456	36.2	258.6	1599	33.1	284.0	3055	34.5	542.6
White	2245	55.8	40.3	2782	57.6	49.9	5027	56.8	90.2
Hispanic	233	5.8	77.4	348	7.2	115.7	581	6.6	193.1
Asian	27	0.7	15.4	26	0.5	14.9	53	0.6	30.3
Other	64	1.6	36.6	71	1.5	40.6	135	1.5	77.1
Total	4025	100.0	63.8	4826	100.0	76.4	8851	100.0	140.2

Prevalence of HIV/AIDS by Race/Ethnicity and Sex

Given the large differences between the racial and ethnic groups as well as the sex of diagnosed clients, this profile will take a closer look at the distribution of race and ethnicity by sex.

Figure 20: Prevalence Rates of HIV/AIDS by Race/Ethnicity and Sex, 2007



The highest prevalence rates for HIV/AIDS are found for males among all racial and ethnic population groups. Among the diagnosed male population, Black males continue to be disproportionately represented. Their prevalence rate is five times the rate of White

males, and almost three times the Hispanic male prevalence rate (Figure 20). In absolute numbers Black men are roughly half the number of their White counterparts. The current rates for both males and females are comparable, but slightly higher, to the rates from the previous year.

A similar picture emerges when comparing the female prevalence rates among the racial/ethnic groups. The highest prevalence rates are among Black females. Their HIV/AIDS prevalence rate is more than ten times higher than their White counterparts and still almost four times the rate of Hispanic females. In absolute numbers among the female diagnosed population, half are Black, while a comparable number of diagnosed females are White. Table 11 shows the absolute numbers, percentages and rates per 100,000 by race/ethnicity and sex.

Table 11: Prevalence Numbers, Percentages and Rates of HIV/AIDS by Race/Ethnicity and Sex in 2007

Race/ Ethnicity	Male			Female		
	Number	%	Rate	Number	%	Rate
Black	2182	30.5	808.1	873	51.2	297.9
White	4340	60.7	157.7	687	40.3	24.3
Hispanic	479	6.7	292.0	102	6.0	74.6
Asian	40	0.6	45.5	13	0.8	14.9
Other	104	1.5	118.3	31	1.8	35.6
Total	7145	100.0	229.7	1706	100.0	53.3

Prevalence of HIV/AIDS by Mode of Transmission

Modes of transmission of the virus were first classified and introduced by the Centers for Disease Control and Prevention (CDC). Those transmission categories are Men having Sex with Men (MSM), Injection Drug Users (IDU), Men having Sex with Men and Injection Drug Users (MSM/IDU), Heterosexual Contact, MSM and Heterosexual Contact, Heterosexual Contact and IDU, and Other. The *Other* category was created to encompass risk categories such as hemophilia and coagulation disorders, transfusion of blood or blood components or tissue transplants, diagnosed mothers, no reported risk mode of transmission, or other categories. Due to the small numbers of all those categories, they are grouped into one category.

During each test for HIV, a person reports information about his or her behavior and events which in turn allows for a risk category classification. In case a person falls into multiple risk categories, the priority follows the sequence of transmission modes as outlined above.

The differences between the transmission mode prevalence rates are considerable. The overwhelming majority of HIV transmissions occurred through Men having Sex with Men (MSM). Tables 12a-b shows the prevalence rates for HIV/AIDS by mode of transmission.

Table 12a: Prevalence Rates (Percents) of Males Living with HIV/AIDS per 100,000 Male Population: Mode of Transmission, 2005-2007

Mode	Rate 2007	Rate 2006	Rate 2005
MSM	140.6 (49.4%)	123.0 (49.6%)	132.4 (51.9%)
MSM/IDU	13.5 (4.7%)	14.3 (5.8%)	14.3 (5.6%)
Total Male Pop.*	229.7 (100.0%)	200.5 (100.0%)	206.7 (100.0%)

*Total Male Pop. = Total Male HIV/AIDS Living Population in Indiana

Table 12b: Prevalence Rates (Percents) of Total Persons Living with HIV/AIDS per 100,000 Population: Mode of Transmission, 2005-2007

Mode	Rate 2007	Rate 2006	Rate 2005
IDU	6.6 (4.7%)	10.4 (8.5%)	11.5 (9.2%)
Heterosexual	23.8 (17.0%)	20.7 (17.0%)	20.6 (16.4%)
MSM/ Heterosexual	10.7 (3.8%)	NA	NA
IDU/ Heterosexual	5.4 (3.9%)	NA	NA
Other	23.2 (16.5%)	18.6 (15.2%)	21.1 (16.8%)
Total Pop.*	140.2 (100.0%)	121.9 (100.0%)	125.3 (100.0%)


*Total Pop. = Total Overall HIV/AIDS Living Population in Indiana

The calculation of the risk category rates differ slightly from the regular rate calculations.

For example, the rate for MSM was calculated by dividing the number of HIV/AIDS cases in Indiana by the number of all men living in Indiana at that time and by multiplying that by 100,000. Given that only males are potentially possible to get diagnosed as MSM (Men Having Sex with Men), the rate calculation is therefore based on only the male part of the general population. The denominator for the calculation of the IDU rate needs to be the entire population of Indiana, since persons of both gender have the potential of becoming IDUs. The rate for MSM/IDU was again calculated with only the male population of Indiana, while Heterosexual Contact and Other included the entire population.

Not all racial and ethnic groups contribute to the risk groups according to their relative size of the general population. Figure 21 breaks out the prevalence rates for transmission modes by race and ethnicity.

Figure 21: Prevalence Rates for HIV/AIDS by Mode of Transmission and Race/Ethnicity, 2007



Consistent across all racial/ethnic categories, the highest HIV/AIDS prevalence rates are associated with MSM. The prevalence rate is especially high for Black MSM, even though *Heterosexual Contacts* and *Other* risk factors register prominently for Blacks as well. Table 12c-d lists the prevalence rates for all racial and ethnic groups.

Table 12c: Prevalence Numbers and Rates of Males Living with HIV/AIDS per 100,000 Male Population: Mode of Transmission and Race/Ethnicity, 2007

Mode of Transmission	Black		Hispanic		White		Asian		Other	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
MSM	1024	379.2	216	131.7	3063	111.3	20	22.7	50	56.9
MSM/IDU	131	48.5	23	14.0	257	9.3	1	1.1	8	9.1
Total Male Pop.*	2,182	808.1	479	292.0	4,340	157.7	40	45.5	104	118.3

*Total Male Pop. = Total Male HIV/AIDS Living Population in Indiana

Table 12d: Prevalence Numbers and Rates of Total Living with HIV/AIDS per 100,000 Population: Mode of Transmission and Race/Ethnicity, 2007

Mode of Transmission	Black		Hispanic		White		Asian		Other	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
IDU	200	35.5	43	14.3	165	3.0	1	0.0	8	4.6
Heterosexual Contact	738	131.1	132	43.9	586	10.5	16	9.1	32	18.3
MSM/Heterosexual	112	41.5	31	18.9	186	6.8	0	0.0	4	4.5
IDU/Heterosexual	150	26.6	16	5.3	168	3.0	0	0.0	7	4.0
Other/No Risk Identified	700	124.3	120	39.9	602	10.8	15	8.6	26	14.9
Total*	3,055	542.6	581	193.1	5,027	90.2	53	30.3	135	77.1


*Total Pop. = Total HIV/AIDS Living Population in Indiana

Prevalence of HIV/AIDS by Health Regions

So far this profile has found that HIV/AIDS is most prevalent among minority men, mainly Blacks in their thirties and forties, whose main risk category is MSM. A look at the regional distribution of the diagnosed population will provide further insight.

Figure 22 shows the prevalence rates per 100,000 people of the population of HIV/AIDS by Indiana’s Health Regions. The rates per health region were calculated using the population estimates 2006.

Figure 22: Prevalence Rates for HIV/AIDS by Health Region, 2007




Indiana shows very distinct regional differences in its prevalence rate for HIV/AIDS. Health Region 6, which covers Indianapolis and the surrounding counties, shows the highest prevalence rate of the entire state, with a rate of 254.8 per 100,000 persons. The next highest rates are in Regions 1 and 7 which includes Lake, Porter and La Porte Counties in northern Indiana and southern counties such as Evansville and Terre Haute in Region 7, with a rate of 167.6 and 103.3 per 100,000 people respectively.

Similar to the Indianapolis metro area, the proximity of the Chicago metro area is influencing the number of diagnosed persons in that region. The highest prevalence rates are all associated with Indiana’s larger cities, such as Muncie/Anderson in Region 5 and Bloomington in south-central Indiana in Region 8.

Figure 23 breaks out the prevalence rate by HIV and AIDS separately for each Health Region. Most of the Health Regions show no large differences between their HIV and AIDS prevalence rates.

Figure 23: Prevalence Rates for HIV and AIDS by Health Region, 2007



The corresponding rates for HIV and AIDS by Health Region are listed in Table 13.

Table 13: Prevalence Numbers, Percentages and Rates per 100,000 Population for HIV, AIDS and HIV/AIDS by Health Region, 2007

Region	HIV			AIDS			HIV/AIDS		
	Number	%	Rate	Number	%	Rate	Number	%	Rate
1	618	15.4	80.8	664	13.8	86.8	1,282	14.5	167.6
2	278	6.9	48.8	309	6.4	54.2	587	6.6	103.0
3	269	6.7	35.8	315	6.5	42.0	584	6.6	77.8
4	94	2.3	26.5	104	2.2	29.3	198	2.2	55.7
5	228	5.7	41.2	313	6.5	56.6	541	6.1	97.8
6	1,856	46.1	115.0	2,256	46.7	139.8	4,112	46.5	254.8
7	326	8.1	45.4	416	8.6	57.9	742	8.4	103.3
8	125	3.1	44.4	131	2.7	46.6	256	2.9	91.0
9	54	1.3	18.2	92	1.9	31.0	146	1.6	49.2
10	177	4.4	43.3	225	4.7	55.0	402	4.5	98.3
Unknown	0	0.0	0.0	1	0.0	0.0	1	0.0	0.0
Total	4025	100.0	63.8	4826	100.0	76.4	8851	100.0	140.2

Please note that in order to calculate the rate for each region, the number of HIV positive and AIDS diagnosed persons is divided by the total number of people living in each region and multiplied by 100,000.

Prevalence of HIV/AIDS by Current State of Residence

At the time of this report, the vast majority of diagnosed persons that are eligible for the programs and services provided by ISDH also reside in the state. Some of the diagnosed persons that had been diagnosed with HIV/AIDS in Indiana have either moved out of the state since their diagnosis, or they lived outside the state of Indiana and were only diagnosed here. Table 14 lists, in descending order, the states of residence of diagnosed persons.

Table 14: Number of Current Indiana Residents with HIV/AIDS by State of Diagnosis, 2007

State of Residence	Number of Diagnosed	State of Residence	Number of Diagnosed
Indiana	8,373	Kansas	5
Florida	61	New York	4
Illinois	54	New Mexico	4
Kentucky	43	Maryland	4
Texas	31	North Carolina	3
Michigan	31	Utah	3
Ohio	27	Foreign Country	3
California	26	Hawaii	3
Arizona	16	Oklahoma	3
Tennessee	14	Mississippi	3
Georgia	13	West Virginia	3
Wisconsin	13	District of Columbia	2
Pennsylvania	11	Rhode Island	2
Minnesota	11	Nebraska	2
Missouri	9	New Jersey	2
Colorado	8	North Dakota	2
Nevada	8	Wyoming	1
Alabama	8	Maine	1
Iowa	8	Connecticut	1
Washington	8	Arkansas	1
Virginia	7	Montana	1
Massachusetts	7	Oregon	1
South Carolina	5		
Louisiana	5	Total	8,851

Within the state of Indiana most diagnosed persons resided in Marion County. Table 15 lists the number of diagnosed persons by Indiana counties, ranked in descending order.

Table 15: Number and Rate per 100,000 Population of Diagnosed Persons with HIV/AIDS by Indiana County of Residence at Time of Report, 2007

County of Residence	Number of Diagnosed	HIV/AIDS Rate	County of Residence	Number of Diagnosed	HIV/AIDS Rate
Marion	3,647	421.4	Daviess	16	52.9
Lake	980	198.3	Perry	16	84.9
Allen	431	124.1	Shelby	16	67.5
St. Joseph	423	2,053.8	Clay	15	55.5
Vanderburgh	265	152.9	Scott	15	5.6
Madison	186	142.4	Wabash	15	44.7
Vigo	182	176.7	Steuben	14	41.6
LaPorte	180	162.9	Dearborn	13	52.1
Clark	177	170.9	Jay	13	60.2
Monroe	173	141.1	Brown	12	79.6
Hamilton	142	56.6	Dubois	12	29.1
Elkhart	132	66.6	Marshall	12	25.4
Porter	122	76.2	Noble	12	25.0
Hendricks	114	86.9	Posey	12	44.8
Delaware	108	94.0	Randolph	12	45.1
Tippecanoe	108	69.2	Vermillion	12	72.1
Johnson	100	75.0	Wells	10	35.5
Floyd	93	128.2	Adams	9	26.7
Howard	90	106.5	Fulton	9	43.6
Wayne	71	103.1	Greene	9	27.0
Putnam	58	156.9	Jasper	9	27.9
Grant	49	70.2	Starke	9	39.0
Miami	45	126.6	Blackford	8	58.8
Bartholomew	44	59.1	Decatur	8	19.1
Kosciusko	40	52.3	Jennings	8	28.1
Morgan	35	49.8	LaGrange	8	21.5
Henry	34	72.4	Spencer	7	15.9
Hancock	32	49.2	Switzerland	7	72.0
Cass	29	72.7	Whitley	7	21.5
Harrison	28	75.7	Carroll	6	29.2
Knox	27	70.6	Fountain	6	34.3
Lawrence	27	58.2	Martin	6	58.0
Boone	26	48.6	Newton	6	42.0
Jackson	26	61.3	Fayette	5	20.3
Warrick	24	42.0	Franklin	4	17.1
Washington	23	82.0	Orange	4	20.3
Montgomery	22	57.6	Ripley	4	14.4
Parke	22	129.3	Rush	4	22.6
Sullivan	22	102.1	Benton	3	33.1
De Kalb	21	42.3	Crawford	2	18.0
Clinton	20	58.5	Pike	2	15.6
Jefferson	19	58.2	Pulaski	2	14.4
Gibson	18	53.9	Union	2	27.4
White	18	73.8	Ohio	1	17.2
Huntington	17	44.7	Tipton	1	6.1
Owen	17	74.8	Warren	0	0.0
			Unknown	1	0.0
			Total	8,851	140.2


New Diagnosis of HIV/AIDS in Indiana

New Diagnosis describes the number and rates of new cases of a disease in a population in a certain amount of time, usually a year. In the case of this report, new diagnosis describes the number of new cases of HIV/AIDS that were treated or reported in Indiana between January 1, 2007 and December 31, 2007 and that were reported in the HIV/AIDS Surveillance Report.

New Diagnosis Rates for Indiana 2007

Indiana started collecting data on HIV and AIDS diagnoses in 1982. Figure 24 shows the development of HIV/AIDS and AIDS over more than two decades, from 1982 up until the end of 2007.

Figure 24: New Diagnosis Numbers of AIDS and HIV/AIDS for Indiana, 1982 to 2007




In the first decade after the recording of diagnosed persons in Indiana began, the numbers steadily climbed, until they reached a peak in the mid-nineties (1996). At that point, the availability and effectiveness of antiviral drugs that, at least temporarily, slowed the progression from HIV infection to AIDS, as well as educational campaigns to stop the spread of the virus brought the rise in the number of diagnosed persons to a halt and, in fact, reversed them for the next four to five years.

Beginning in or around the year 2000, however, the number of new HIV/AIDS and AIDS diagnoses started to plateau and did not change much over the course of three years. The numbers did rise in 2006, however, and slightly decrease in 2007. Delays in reporting may affect this slight decrease.

In 2007, the number of newly diagnosed persons with HIV/AIDS was 509, a fairly small decrease from 2006 (540). The new diagnosis rate is calculated at 8.1 per 100,000 persons, compared to a new diagnosis rate of 8.7 in 2006.

Similar to the prevalence numbers, the group of newly diagnosed persons is predominantly male.

Figure 25: New Diagnosis Rates for HIV/AIDS by Sex, 2007



Males continue to have a more than three times higher new diagnosis rate than females. They contributed more than three quarters to the new diagnosis rate in 2007.

A more detailed look at the new diagnosis rates for HIV and AIDS separately is provided in Figure 26, which shows the difference in new diagnosis rates by sex for HIV and AIDS separately.

Figure 26: New Diagnosis Rates for HIV and AIDS by Sex, 2007




Table 16 shows the absolute number for HIV, AIDS and HIV/AIDS New Diagnosis in Indiana for 2007.


Table 16: New Diagnosis Numbers, Percentages and Rates per 100,000 Population for HIV, AIDS, and HIV/AIDS by Sex, 2007

Sex	HIV			AIDS			HIV/AIDS		
	Number	%	Rate	Number	%	Rate	Number	%	Rate
Male	249	74.8	8.0	140	79.5	4.5	389	76.4	12.5
Female	84	25.2	2.6	36	20.5	1.1	120	23.6	3.7
Total	333	100.0	5.3	176	100.0	2.8	509	100.0	8.1

New Diagnosis of HIV/AIDS by Age


In 2007, the new diagnosis rate for HIV/AIDS peaked among 30 to 39 year olds (17.6 per 100,000).

Figure 27: New Diagnosis Rates for HIV/AIDS by Age, 2007



A more detailed look at the HIV and AIDS new diagnosis rates by age for 2007 reveal a more detailed picture of the age distribution among newly diagnosed persons.

Figure 28: New Diagnosis Rates for HIV and AIDS by Age, 2007



For most all age groups the HIV new diagnosis rate exceeds the rate of newly diagnosed AIDS cases. The vast majority of newly diagnosed HIV cases in 2007 were in the age ranges of 20 to 24, while the majority of new AIDS diagnoses occurred for persons in the age ranges of 40 to 49 years of age.

Table 17 shows the absolute numbers, percentages and rates for the combined disease as well as the separate diagnoses for 2007.


Table 17: New Diagnosis Numbers, Percentages and Rates for HIV, AIDS, and HIV/AIDS by Age, 2007

Age	HIV			AIDS			HIV/AIDS		
	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000	Number	%	Rate/ 100,000
0 to 9	<5	NA	NA	<5	NA	NA	<5	NA	NA
10 to 14	<5	NA	NA	<5	NA	NA	<5	NA	NA
15 to 19	18	5.4	4.0	4	2.3	0.9	22	4.3	4.9
20 to 24	58	17.4	13.1	12	6.8	2.7	70	13.8	15.9
25 to 29	57	17.1	12.8	19	10.8	4.3	76	14.9	17.1
30 to 39	99	29.7	11.9	48	27.3	5.7	147	28.9	17.6
40 to 49	63	18.9	6.6	67	38.1	7.1	130	25.5	13.7
50 to 59	26	7.8	3.1	23	13.1	2.8	49	9.6	5.9
60 to 64	7	2.1	2.5	<5	NA	NA	9	1.8	3.2
over 65	<5	NA	NA	<5	NA	NA	<5	NA	NA
Total	333	100.0	5.3	176	100.0	2.8	509	100.0	8.1

New Diagnosis Rate of HIV/AIDS by Race/Ethnicity

In Figure 29 the new diagnosis rates are shown by race and ethnicity. In order to calculate the rate per 100,000 persons, the number of diagnosed persons for each race and ethnicity was divided by the number of the entire Indiana population that were identified in the 2006 Census Estimates as belonging to that particular racial and ethnic category.

Figure 29: New Diagnosis Rates of HIV/AIDS by Race/Ethnicity, 2007



The overwhelming majority of newly diagnosed persons were Black. They had a new diagnosis rate of 37.1 per 100,000 people of the population, nearly twice as large as the next largest group of Hispanics (14.3 per 100,000). It is interesting to note that Blacks make up a minority of the general population of about 8.9%, yet they account for 41.1% of all new cases of HIV/AIDS in 2007.


The New Diagnosis numbers for HIV/AIDS for absolute numbers, percentages and rates per 100,000 are listed in Table 18.

Table 18: New Diagnosis Numbers, Percentages and Rates per 100,000 Population for HIV/AIDS by Race/Ethnicity, 2007

	Number	Percent	Rate
Black	209	41.1	37.1
White	236	46.4	4.2
Hispanic	43	8.4	14.3
Asian	9	1.8	5.1
Other	12	2.4	6.9
Total	509	100.0	8.1

The absolute numbers for the combined disease as well as the percentage numbers show that all Black and Hispanic groups are over-represented in the number of newly diagnosed persons, when compared to their part of the overall population.

Figure 30: New Diagnosis Rates for HIV and AIDS by Race/Ethnicity, 2007



The separate view of HIV and AIDS by race and ethnicity reveals further information about the different behavior of new diagnoses. Displayed in Figure 32 are the new HIV diagnosis and newly diagnosed cases of AIDS by race and ethnicity. Blacks show the highest rate of new HIV diagnosis as well as new AIDS diagnoses among all racial and ethnic groups in Indiana. The

second largest minority population group, Hispanics, have also the second largest new diagnosis rates for both HIV and AIDS. In other words, the results in Figure 29 and Figure 30 show that new diagnosis of HIV and AIDS are spreading much more rapidly among minority population groups than among Whites, a finding that is consistent with the status of the prevalence ratings. However, by absolute numbers, the new diagnosis for Whites outnumbers all other racial and ethnic groups.


Table 19 lists the absolute numbers of newly diagnosed persons by race/ethnicity, as well as the percentage of the overall diagnosis and the rates per 100,000 people of the population.

Table 19: New Diagnosis Numbers, Percentages and Rates per 100,000 Population for HIV and AIDS by Race/Ethnicity, 2007

Race/ Ethnicity	HIV			AIDS		
	Total	Percent	Rate	Total	Percent	Rate
Black	148	44.4	26.3	61	34.7	10.8
White	145	43.5	2.6	91	51.7	1.6
Hispanic	24	7.2	8.0	19	10.8	6.3
Asian	6	1.8	3.4	<5	NA	NA
Other	10	3.0	5.7	<5	NA	NA
Total	333	100.0	5.3	176	100.0	2.7

In addition to racial and ethnic differences in the new diagnosis rates, there are also differences in the number of male and female new diagnosis among racial and ethnic groups. Figure 31 shows the new diagnosis rate breakout by race/ethnicity and sex.

Figure 31: New Diagnosis Rates for HIV/AIDS by Race/Ethnicity and Sex, 2007



The new diagnosis rate results for males and females mirror the earlier assessments, in which new diagnoses rates are highest among the male groups of racial and ethnic minorities. The rates for Black and Hispanic males are three to six times the White male new diagnosis rate (see Figure 31 and Table 19).

Comparing the female new diagnosis rates among these racial/ethnic groups shows a similar result. HIV/AIDS new diagnosis rates are lowest among White females and highest among Black females. In absolute numbers, Black females make up the majority of new diagnosis with HIV. The results for all racial and ethnic groups for the combined disease HIV/AIDS are summarized in Table 20.

Table 20: New Diagnosis Numbers, Rates per 100,000 Population and Percentages for HIV/AIDS by Race/Ethnicity and Sex, 2007

Race/ Ethnicity	Male			Female		
	Total	Rate	Percent	Total	Rate	Percent
Black	154	39.6	57.0	55	45.8	18.8
White	187	48.1	6.8	49	40.8	1.7
Hispanic	34	8.7	20.7	9	7.5	6.6
Other	7	1.8	8.0	5	4.2	5.7
Asian	7	1.8	8.0	<5	NA	NA
Total	389	100.0	12.5	120	100.0	3.7

The rates in Table 20 were calculated by dividing the absolute number of new cases of HIV/AIDS by the number of the racial and ethnic male or female population respectively, and multiplying that number by 100,000. The reduction of the absolute numbers to the rates per 100,000 allows for direct comparison of rates between the different racial and ethnic groups as well as between the sex categories.

New Diagnosis of HIV/AIDS by Mode of Transmission

The New Diagnosis rates of HIV/AIDS vary widely by mode of transmission for 2005-2007, as shown in Table 21a-b.

Table 21a: New Diagnosis Rates (Percents) of Males Reported with HIV/AIDS per 100,000 Male Population: Mode of Transmission, 2005-2007

Mode	Rate 2007	Rate 2006	Rate 2005
MSM	6.7 (40.7%)	8.0 (45.4%)	5.9 (42.4%)
MSM/IDU	0.3 (1.8%)	0.3 (3.0%)	0.7 (5.0%)
Total Male Pop.*	12.5 (100.0%)	13.4 (100.0%)	10.8 (100.0%)

*Total Male Pop. = Total Male HIV/AIDS New Diagnosis in Indiana

Table 21b: New Diagnosis Rates (Percents) of Total Persons Reported with HIV/AIDS per 100,000 Population: Mode of Transmission, 2005-2007

Mode	Rate 2007	Rate 2006	Rate 2005
IDU	0.2 (2.6%)	0.2 (1.9%)	0.2 (3.3%)
Heterosexual	1.3 (15.5%)	1.1 (12.6%)	1.0 (14.7%)
MSM/ Heterosexual	0.3 (2.0%)	NA	NA
IDU/ Heterosexual	0.2 (2.0%)	NA	NA
Other	2.9 (35.6%)	2.8 (32.6%)	2.4 (34.6%)
Total Pop.*	8.1 (100.0%)	8.7 (100.0%)	6.8 (100.0%)

*Total Pop. = Total HIV/AIDS New Diagnosis in Indiana


NA= Not Applicable

Note: For categories MSM and MSM/IDU rates are relative to the number of men, not the overall population. Thus, comparisons across transmissions should not be made.

The vast majority of new cases registered in 2007 are in the category of Men having Sexual Contact with Men (MSM). The *MSM* rate decreased from the previous year (8.7/100,000).

In Figure 32, HIV/AIDS New Diagnosis rates are computed separately by race/ethnicity categories and mode of transmission.

Figure 32: New Diagnosis Rates of HIV/AIDS by Mode of Transmission and Race/Ethnicity, 2007



Note: For categories MSM, and MSM/IDU rates are relative to the number of men Not the overall population. Thus, comparisons across transmissions should not be made. However, comparisons by Race/Ethnicity for each transmission category can be made.

Consistent across all race/ethnic categories, the highest HIV/AIDS new diagnosis rates are associated with *MSM*. For nearly all race/ethnic categories, the *Heterosexual contact* risk category accounts for the second highest HIV/AIDS New Diagnosis rates. The new diagnosis numbers and rates per 100,000 for all racial and ethnic groups by mode of transmission are listed in Tables 22a-b.

Table 22a: New Diagnosis Numbers and Rates of Males Reported with HIV/AIDS per 100,000 Male Population: Mode of Transmission and Race/Ethnicity, 2007

Mode of Transmission	Black		Hispanic		White		Other	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate
MSM	69	25.6	16	9.8	113	4.1	6	6.8
MSM/IDU	<5	NA	<5	NA	5	0.2	<5	NA
Total Male Pop.*	154	57.0	34	20.7	187	6.8	14	15.9

*Total Male Pop. = Total Male HIV/AIDS Newly Diagnosed in Indiana
NA= Not Applicable

Table 22b: New Diagnosis Numbers and Rates of Total Reported with HIV/AIDS per 100,000 Population: Mode of Transmission and Race/Ethnicity, 2007

Mode of Transmission	Black		Hispanic		White		Other	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate
IDU	5	0.9	<5	NA	6	0.1	<5	NA
Heterosexual Contact	40	7.1	10	3.3	24	0.4	<5	NA
MSM/Heterosexual	6	2.2	<5	NA	<5	NA	<5	NA
IDU/Heterosexual	5	0.9	<5	NA	5	0.1	<5	NA
Other/No Risk Identified	82	14.6	15	5.0	79	1.4	<5	NA
Total*	209	37.1	43	14.3	236	4.2	21	12.0


*Total Pop. = Total HIV/AIDS Newly Diagnosed in Indiana
NA= Not Applicable

In order to calculate the rate/100,000 people of the general population, the absolute number of people per risk category was divided by the number of the corresponding general population. However, the rate for MSM and MSM/IDU was calculated using the corresponding number of males of that particular racial and ethnic group.

New Diagnosis Rate for HIV/AIDS by Health Regions and Counties

The geographic distribution of new diagnosis cases in Indiana shows regional differences. For the most part the highest new diagnosis rates are corresponding to the population size and proximity of the health region to large urban areas in the state. The distribution of new diagnosis rates for all ten Indiana Health Regions is provided in Figure 33.

Figure 33: New Diagnosis Rates of HIV/AIDS by Health Region in Indiana, 2007



Region 6, which corresponds to the Greater Indianapolis area, shows by far the largest share in the number of newly diagnosed people. Their rate of 14.4 persons per 100,000 people of the population far exceeds the rates of the next closest regions (1, 8, and 10), which correspond to the areas around Indiana's northern and southern regions respectively.


The New Diagnosis details of the combined disease HIV/AIDS for Indiana Health Regions are listed in Table 23.

Table 23: New Diagnosis Numbers, Percentages and Rates per 100,000 Population for HIV/AIDS by Health Region, 2007

Health Region	HIV/AIDS		
	Total	Percent	Rate
1	93	18.3	12.2
2	25	4.9	4.4
3	31	6.1	4.1
4	11	2.2	3.1
5	20	3.9	3.6
6	232	45.6	14.4
7	32	6.3	4.5
8	14	2.8	5.0
9	10	2.0	3.4
10	33	6.5	8.1
Unknown	8	1.6	0.1
Total	509	100.0	8.1

The New Diagnosis rate distribution with HIV and AIDS for each Health Region is displayed in Figure 34 below.

Figure 34: New Diagnosis Rates for HIV and AIDS by Health Region, 2007



Consistent with the numbers for the combined disease the majority of newly diagnosed cases of HIV and AIDS occur in Region 6, the greater Indianapolis area.

The separate HIV and AIDS New Diagnosis numbers are listed in Table 24 by Health Region.

Table 24: New Diagnosis Numbers, Percentages and Rates per 100,000 Population for HIV and AIDS by Health Region, 2007

Health Region	HIV			AIDS		
	Number	Percent	Rate	Number	Percent	Rate
1	56	16.8	7.3	37	21.0	4.8
2	21	6.2	3.7	<5	NA	NA
3	18	5.3	2.4	13	7.7	1.7
4	5	1.5	1.4	6	3.4	1.7
5	14	4.2	2.5	6	3.4	1.1
6	166	49.8	10.3	66	37.5	4.1
7	21	6.3	2.9	11	6.3	1.5
8	10	3.0	3.6	<5	NA	NA
9	5	1.5	1.7	5	2.8	1.7
10	21	6.3	5.1	12	6.8	2.9
Unknown	4	1.2	0.1	4	2.4	0.1
Total	341	100.0	5.4	168	100.0	2.7

In order to refine the geographic distribution of the newly diagnosed cases this profile also takes a look at the number of cases per county. Table 25 below is listing the new diagnosis numbers for HIV/AIDS by the counties of residence, in declining order of magnitude for the combined disease. For reasons of confidentiality, no new diagnosis numbers smaller than 5 is reported. All counties with fewer than five diagnosed persons are combined into an *Other* category.

Table 25: New Diagnosis Numbers for HIV/AIDS by County, 2007

County	HIV/AIDS Number
Marion	201
Lake	74
Allen	22
St. Joseph	20
Clark	14
Hendricks	13
LaPorte	12
Monroe	10
Floyd	9
Porter	8
Vanderburgh	8
Hamilton	7
Madison	7
Vigo	7
Elkhart	6
Putnam	5
Johnson	5
Other	81
Total	509


Pediatric Classification and HIV Status of Mothers

All infants born to an HIV positive mother should be reported to the state health department, even though the final HIV status of the child is not known until later. By the end of 2007, a total of 642 children had been born to HIV positive mothers since the beginning of record keeping. During the year 2007, two new cases of pediatric infection was recorded. All children recorded in the eHARS database are classified in one of four categories:

- **Exposed:** Children that are born to HIV+ women, but their laboratory testing has not yet determined their HIV status
- **HIV:** Children that are born to HIV+ women and their laboratory has confirmed their HIV+ status
- **AIDS:** Children that are born to HIV+ women and they meet the definition for pediatric AIDS
- **Seroreverter:** Definitely Not Diagnosed, the laboratory testing has confirmed that child is definitely not diagnosed
- **DNI:** Definitely Not Infected

In Figure 35 the distribution of children among these four categories is shown. The numbers are cumulative.

Figure 35: Cumulative Number of Children born to HIV positive Mothers up until December 31, 2007



Note: A total of 24 cases did not reveal status at the time of this report.

Table 26 shows the number of children that were born to HIV positive mothers by the time of the mother's diagnosis.

Table 26: Cumulative Numbers and Percentages of Children born to HIV positive Mothers, 2007

HIV Status of Mother	Perinatal Exposure	Pediatric HIV	Pediatric AIDS	Seroreverter	Total	Percent
Refused HIV Testing	0	0	1	0	1	0.2%
HIV+ before Pregnancy	65	8	9	209	291	47.1%
HIV+ during Pregnancy	19	7	5	105	136	22.0%
HIV+ at Delivery	4	2	0	6	12	1.9%
HIV+ sometime before Birth	24	6	3	41	74	12.0%
HIV+ after Birth	2	18	28	30	78	12.6%
HIV+ Time Unknown	3	4	5	4	16	2.6%
Unknown	0	2	6	2	10	1.6%
Total	117	47	57	397	618	100%


Note: A total of 24 cases did not reveal status at the time of this report.

The majority of children were born to mothers whose HIV positive status was determined either before (47.1%) or during (22.0%) a pregnancy. The time of detection of the HIV positive status of the mother is important in estimating the risk to the children of diagnosed mothers. Early detection of the mother's status improves the chances of preventing the spread of the virus from the mother to the child, either during birth or after the child is born. Accordingly, the number of children that were diagnosed as HIV positive or meeting the criteria for pediatric AIDS was about two times larger if the status of the mother was diagnosed as HIV positive after the child was born, or where the time of infection in regards to the diagnosis was unknown.

Two new pediatric case of HIV or AIDS was diagnosed in 2007.

The ethnic and racial distribution of the group of children that were born to diagnosed mothers is presented in Figure 36. The figure includes those children that are currently in the *Exposed* category, pending the outcome of their laboratory results, as well as those that were diagnosed as either *HIV positive*, *AIDS*, or *Definitely Not Diagnosed*.

Figure 36: Cumulative Percentages of Children born to HIV+ Mothers by Race/Ethnicity, 2007



More than half of all children born to diagnosed mothers were Black. Table 27 shows the absolute numbers and corresponding percentages for all four categories by race and ethnicity. The different percentages of children in each category are consistent with the overall distribution of children by category and race/ethnicity.

Table 27: Cumulative Numbers and Percentages of Children born to HIV+ Mothers by Race/Ethnicity, 2007

Race/ Ethnicity	Exposure	%	Pediatric HIV	%	Pediatric AIDS	%	DNI	%	Total	%
Black	82	68.9	26	48.1	29	36.7	206	51.9	343	52.9
White	21	17.6	18	38.	40	55.7	153	38.5	232	36.8
Hispanic	12	10.1	4	7.4	2	2.5	29	7.3	47	7.2
Other	4	3.4	3	5.6	4	5.1	9	2.3	20	3.1
Total	119	100.0	51	100.0	75	100.0	397	100.0	642	100.0

Indiana law requires the primary prenatal care provider to offer the pregnant women HIV information, counseling and voluntary testing. Medical studies have shown that pregnant women who are HIV positive can reduce the risk of passing the virus on to their children by two-thirds with proper perinatal care and antiviral treatment during pregnancy, labor, delivery and to the

child after birth. Table 28 shows the number of children by their infection status broken out by the time the mother received drug treatment to lower her viral load.

Table 28: Cumulative Number of Children by their Infection Status and by the Availability of Drugs, 2007

Mother received drug...	...prior to Pregnancy	...during Pregnancy	...during Delivery	Child received drugs
Exposure	30	72	78	93
Pediatric HIV	1	10	11	15
Pediatric AIDS	0	5	0	3
DNI	100	250	250	294

In case of the mother receiving antiviral drugs before the pregnancy, no case of pediatric HIV or AIDS has been diagnosed. When taking the drugs during pregnancy the number of diagnosed children is still very low compared to the not diagnosed children of that same group.

Table 29: Numbers and Percentages of Children by the Mother's Mode of Transmission and by the Child's Infection Status, 2007

Mode of Transmission of Mother	Perinatal Exposure	Pediatric HIV	Pediatric AIDS	Seroreverter	Total	Percent
IDU	10	11	13	46	80	14.7%
Sexual contact with IDU	17	9	16	77	119	21.8%
Sexual contact with Bisexual Male	9	3	5	38	55	10.1%
Sexual contact with Male with Hemophilia	1	0	0	8	9	1.7%
Sexual contact with HIV+ Transfusion Recipient	1	0	0	2	3	0.6%
Sexual Contact with Male with HIV or AIDS	35	18	20	198	271	49.7%
Received Blood Transfusion	0	2	2	4	8	1.5%
Total	73	43	56	373	545*	100.0%

*Does not equal the total number of pediatric cases because some cases did not reveal the mode of transmission, and some cases include more than one mode of transmission.


The majority of mothers (49.7%) were exposed to HIV or AIDS through sexual contact with a HIV+ male. Another 21.8% of mothers were exposed due to IDU, either by using personally using intravenous drugs or through sexual contact with someone who was an IDU.

Department of Corrections

The Indiana Department of Corrections (DOC) started conducting mandatory HIV tests as part of the general intake process for every inmate at the beginning of their corrections term on July 1, 2002. Currently, no such test is administered when an inmate is released.

There are currently 352 inmates in Indiana Department of Correction facilities. The vast majority are male (316 or 89.8%). The majority of HIV-positive inmates are Black (59.4%), followed by White (37.5%), and Hispanics (3.1%). The number of those diagnosed remained virtually the same from the previous year.


Figure 37: Number of Inmates with HIV/AIDS by Race/Ethnicity, 2007



Source: ISDH 2007

The risk category distribution for the inmates is shown in Figure 38.

Figure 38: Number of Inmates with HIV/AIDS by Mode of Transmission, 2007




Source: ISDH 2007

The majority of inmates self-identify their risk category as MSM.

The age distribution of diagnosed inmates is shown in Figure 39.

Figure 39: Number of Inmates with HIV/AIDS by Age, 2007



Source: ISDH 2007


The majority of inmates with HIV/AIDS fall between the 30 to 49 age group.

Mortality

There is a difference between the number of deaths of persons with HIV/AIDS and the number of deaths due to HIV or AIDS. Deaths reported by the HIV/AIDS Surveillance program include all deaths of persons who were diagnosed with HIV or diagnosed with AIDS. The deaths reported by Vital Records (death certificates) include only those who died as a result of AIDS and such was identified on the death certificate. The deaths reported here are deaths of persons who were diagnosed with HIV or diagnosed with AIDS regardless of the cause of death. For example, the death may have been due to an automobile accident. Even though the person did not die due to the presence of HIV, the person is no longer living in Indiana and, therefore, not contributing to understanding and planning for HIV prevention or medical services.

Figure 40 shows the number of deaths of persons with HIV/AIDS since 1982. Shown in the figure are the absolute numbers of deaths in 1995 at the peak of annual mortality of diagnosed persons and in 2007. The decline in annual death numbers in 1995 was due to the availability and effectiveness of antiretroviral drugs.

Figure 40: Number of HIV/AIDS Deaths by Year in Indiana, 1982 to 2007



In 2007, 210 persons that were diagnosed with HIV/AIDS died, up from 121 in 2006. That equates to a rate of 2.37 per 100 persons compared to a rate of 1.6 per 100 persons in 2006. The rate continues to increase in 2007 mostly due to increased efforts by surveillance to perform routine death matches.

The mortality rate is calculated by dividing the number of persons that died by the number of the diagnosed population and multiplying that by 100. The trend of recent years of declining mortality numbers was reversed in 2005. Table 30 shows the absolute number, percentages and rates broken out by sex for 2007.


Table 30: Mortality Numbers, Percentages and Rates of HIV/AIDS by Sex, 2007

Sex	Number	Percentage	Rate/100
Male	166	79.0%	1.88
Female	44	21.0%	0.50
Total	210	100.0%	2.37

Diagnosed males were about four times more likely than females to have died in 2007. The mortality rates reflect the gender composition of the diagnosed population, where males have a higher prevalence rate than females do.

Figure 41 shows the breakout of the mortality rate by age groups.

Figure 41: Mortality Rates of HIV/AIDS by Age of Death in Indiana, 2007



The absolute numbers, percentages and rates by age group are listed in Table 31. Also included in Table 31 are the total numbers of diagnosed persons by age group that was used to calculate the mortality rates.

Table 31: Mortality Numbers, Percentages and Rates of HIV/AIDS by Age of Death, 2007

Age Group in Years	Number of Deaths	Percent	Mortality Rate/100	Total Number of Diagnosed Persons
0 to 9	0	0.0%	0.00	0
10 to 14	0	0.0%	0.00	41
15 to 19	1	0.5%	0.00	39
20 to 29	11	5.2%	1.39	789
30 to 39	39	18.6%	1.80	2,169
40 to 49	92	43.8%	2.49	3,700
50 to 59	52	24.8%	3.16	1646
60 to 64	10	4.8%	3.60	278
65+	5	2.4%	2.65	189
Total	210	100.0%	2.37	8,851

Figure 42 presents further detail on the racial and ethnic characteristics of the deceased persons in 2007.

Figure 42: Mortality Rates of HIV/AIDS by Race and Ethnicity, 2007





Table 32 lists the absolute mortality numbers, percentages and rates for the different racial and ethnic groups for 2007.

Table 32: Mortality Numbers, Percentages and Rates of HIV/AIDS by Race/Ethnicity, 2007

Race/Ethnicity	Number	Percent	Rate/100	Number of HIV/AIDS Living Persons by Race/Ethnicity
White	132	62.9%	2.6	5,027
Black	57	27.1%	1.9	3,055
Other	7	3.3%	3.7	188
Hispanic	14	6.7%	2.4	581
Total	210	100.0%	2.4	8,851

The distribution of numbers of transmission modes associated with those persons that died in 2007 in Figure 43 shows mostly a similar picture to the prevalence and new diagnosis rates earlier.

Figure 43: Mortality Rates of HIV/AIDS by Mode of Transmission for Indiana, 2007



Persons identified as IDU, MSM/IDU and IDU/Hetero show to have the highest mortality rates. Those reported as MSM only have the third lowest mortality rate. Table 33 shows the corresponding numbers, rates and percentages for the risk categories.


Table 33: Mortality Numbers, Rates and Percentages of HIV/AIDS by Mode of Transmission, 2007

Mode of Transmission/ Risk Category	Number of Deaths	Rate/100	Percent
MSM	95	2.17	45.2%
IDU	15	3.60	7.1%
Heterosexual Contact	30	1.99	14.3%
MSM/ IDU	13	3.10	6.2%
MSM/ Heterosexual Contact	5	1.50	2.4%
IDU/ Heterosexual Contact	11	3.23	5.2%
No Risk Reported	41	2.80	19.5%
Total	210	2.37	100.0%

*One participant was listed in the “other” category

Finally, the distribution of deaths among the diagnosed population in 2007 in Indiana shows large differences. The mortality rate distribution by Health Region is shown in Figure 44.

Figure 44: Mortality Rates of HIV/AIDS by Health Regions for Indiana, 2007



The mortality rate for each region was calculated by dividing the number of diagnosed people that died in that region by the number of diagnosed people that lived in that region in 2007. The “Unknown” variable includes those with either no county listed or a county outside of Indiana listed.

*** Health Region Key**

Region	Area
1	Northwest Indiana - Lake Region
2	Northcentral Indiana - Elkhart
3	Northeast Indiana - Fort Wayne
4	Westcentral Indiana - Lafayette
5	Eastcentral Indiana - Marion
6	Central Indiana - Indianapolis
7	Southwestern Indiana - Evansville/Terre Haute
8	Bloomington Area
9	Southeastern Indiana - Cincinnati Area
10	Southern Indiana - Louisville Area

The corresponding numbers, percentages and rates by Health Region are in Table 34.

Table 34: Mortality Numbers, Percentages and Rates of HIV/AIDS by Health Regions for Indiana, 2007

Health Region	Number of Deaths	Percent	Rate/100	Number of Diagnosed Persons by Region
1	32	26.4%	2.72	1,177
2	17	14.0%	3.46	492
3	13	10.7%	2.25	578
4	2	1.7%	1.01	198
5	15	12.4%	2.77	541
6	71	58.7%	1.73	4,112
7	22	18.2%	3.01	730
8	8	6.6%	3.13	256
9	9	7.4%	6.16	146
10	15	12.4%	3.80	395
Unknown	6	5.0%	2.65	226
Total	210	173.6%	2.37	8,851

Finally, when ranked among the other states of the U.S., Indiana ranks 25th (up from 33rd in 2003) in the number of HIV related deaths in 2005, the last year that national data was available for comparisons.⁵

⁵ Kaiser Family Foundation, (<http://www.statehealthfacts.kff.org>), 2005

Migration Patterns

By the end of 2007, a total of 478 persons that were diagnosed in Indiana with either HIV or AIDS or HIV/AIDS and were not known to have died, had moved out of the state (Out-Migration) as compared with 268 in 2006. At the same time, 1,455 persons that were diagnosed with either HIV or AIDS or HIV/AIDS in a state other than Indiana, had moved here by the end of 2007 (In-Migration) as compared with 1,308 in 2006. Table 36 lists the cumulative numbers of Out-Migrants by the state to which they moved, while Table 35 lists the cumulative number of In-Migrants by the state in which they were diagnosed with HIV or AIDS.

Table 35: Numbers and Percentages of Persons Diagnosed with HIV or AIDS in Indiana and Currently Living outside the State, 2007 (Out-Migration)


Current State of Residence	Number	Percent	Current State of Residence	Number	Percent
Alabama	8	1.67%	Mississippi	3	0.63%
Arkansas	1	0.21%	Montana	1	0.21%
Arizona	16	3.35%	North Carolina	3	0.63%
California	26	5.44%	North Dakota	2	0.42%
Colorado	8	1.67%	Nebraska	2	0.42%
Connecticut	1	0.21%	New Jersey	2	0.42%
District of Columbia	2	0.42%	New Mexico	4	0.84%
Foreign Country	3	0.63%	Nevada	8	1.67%
Florida	61	12.76%	New York	4	0.84%
Georgia	13	2.72%	Ohio	27	5.65%
Hawaii	3	0.63%	Oklahoma	3	0.63%
Iowa	8	1.67%	Oregon	1	0.21%
Illinois	54	11.30%	Pennsylvania	11	2.30%
Kansas	5	1.05%	Rhode Island	2	0.42%
Kentucky	43	9.00%	South Carolina	5	1.05%
Louisiana	5	1.05%	Tennessee	14	2.93%
Massachusetts	7	1.46%	Texas	31	6.49%
Maryland	4	0.84%	Utah	3	0.63%
Maine	1	0.21%	Virginia	7	1.46%
Michigan	31	6.49%	Washington	8	1.67%
Minnesota	11	2.30%	Wisconsin	13	2.72%
Missouri	9	1.88%	West Virginia	3	0.63%
			Wyoming	1	0.21%
			Total	478	100.0%

Table 36: Numbers and Percentages of Persons that were Diagnosed with HIV or AIDS outside of Indiana and Migrated to Indiana, 2007 (In-Migration)

State of Diagnosis	HIV/AIDS	Percent HIV/AIDS
Alabama	27	1.85%
Alaska	3	0.21%
Arizona	30	2.06%
Arkansas	13	0.89%
California	110	7.55%
Colorado	27	1.85%
Connecticut	6	0.41%
Delaware	4	0.27%
District of Columbia	16	1.10%
Florida	183	12.56%
Foreign Country	7	0.48%
Georgia	40	2.75%
Hawaii	1	0.07%
Illinois	190	13.04%
Iowa	9	0.62%
Kansas	15	1.03%
Kentucky	91	6.25%
Louisiana	26	1.78%
Maine	2	0.14%
Maryland	13	0.89%
Massachusetts	4	0.27%
Michigan	61	4.19%
Minnesota	27	1.85%
Mississippi	27	1.85%
Missouri	49	3.36%
Montana	1	0.07%
Nebraska	2	0.14%
Nevada	20	1.37%
New Hampshire	2	0.14%
New Jersey	24	1.65%
New Mexico	6	0.41%
New York	41	2.81%
North Carolina	17	1.17%
Ohio	89	6.11%
Oklahoma	9	0.62%
Oregon	6	0.41%
Pennsylvania	22	1.51%
Rhode Island	1	0.07%
South Carolina	20	1.51%
Tennessee	37	2.54%
Texas	85	5.83%
Utah	4	0.27%
Vermont	1	0.07%
Virginia	24	1.65%
Washington	19	1.30%
Washington, DC	0	0.00%
West Virginia	5	0.34%
Wisconsin	37	2.54%
Wyoming	2	0.14%
Total	1,455	100.0%

A look at the sex distribution of diagnosed people reveals a large difference between male and female migrant numbers. Figure 45 shows the cumulative numbers of diagnosed persons migrating to and from Indiana by sex. The numbers for both the migration to Indiana and out of the state reflect the total number of diagnosed persons that have been recorded since 1982 up until 2007.


Figure 45: Cumulative Number of Migrants that were Diagnosed with HIV or AIDS by Sex, including 2007



For both migration directions, males outnumber females by four to five times. They make up roughly 84% of the migrants. There is virtually no difference between the gender distribution of diagnosed people that move to Indiana or of those that are leaving the state, after they have been diagnosed here, other than the difference in absolute numbers.

There are, however, differences when considering the racial and ethnic composition of both migrating groups. Figure 46 shows the number of migrating persons that were alive at the time of this report by their race and ethnicity.

Figure 46: Cumulative Number of Migrants that were Diagnosed with HIV or AIDS by Race/Ethnicity and Migration Direction, including 2007




In absolute numbers, the “In-Migrants” outnumber “Out-Migrants”. However, in terms of percentages about the same share of diagnosed persons of minority backgrounds have left the state when compared to those that have moved to Indiana. The same is true for diagnosed White persons. About the same percentage of White and minority background migrants have moved to Indiana when compared to those that have left it. However, in terms of absolute numbers, the number of in-migrants exceeds that of persons leaving the state. In other words, since 1982 Indiana’s population of HIV/AIDS diagnosed people has seen a net growth because of migration. The absolute numbers and corresponding percentages are listed in Table 37.

Table 37: Cumulative Numbers and Percentages of Migrants that were Diagnosed with HIV or AIDS by Race/Ethnicity and Migration Direction, including 2007

Race/Ethnicity	In-Migration	Percent	Out-Migration	Percent
White	824	56.63%	292	61.09%
Black	484	33.26%	156	32.64%
Other	35	2.41%	9	1.88%
Hispanic	112	7.70%	21	4.39%
Total	1,455	100.0%	478	100.0%

The group of migrants that moved to Indiana after they were diagnosed with HIV/AIDS did settle in various parts of the state. Figure 47 shows the distribution of in-migrants by Health Region in Indiana.

Figure 47: Cumulative Percentages of In-Migrants that were Diagnosed with HIV or AIDS by Health Region, including 2007




Region Key:

Region	Area
1	Northwest Indiana - Lake Region
2	Northcentral Indiana - Elkhart
3	Northeast Indiana - Fort Wayne
4	Westcentral Indiana - Lafayette
5	Eastcentral Indiana - Marion
6	Central Indiana - Indianapolis
7	Southwestern Indiana - Evansville/Terre Haute
8	Bloomington Area
9	Southeastern Indiana - Cincinnati Area
10	Southern Indiana - Louisville Area

The Health Regions surrounding the urban centers of the state attracted the largest number of people coming to Indiana. Health Region 6, the greater Indianapolis area, attracted almost four out of ten migrants (39%) of all in-migrants alone.

The distribution of risk categories among migrants is shown in Figure 48.

Figure 48: Number of Migrants that were Diagnosed with HIV or AIDS by Mode of Transmission, including 2007



Similar to Indiana’s resident diagnosed population, *MSM* is the dominant transmission mode among the migrant population. Over half of persons moving to Indiana (52%) were associated with the *MSM* risk category. This ratio is also consistent with the overall diagnosed population, where about 51% were associated with *MSM*. The same is true for *MSM/IDU* and *Heterosexual Contacts*, which both are represented among the migrant population in nearly the same ratios as the current diagnosed population at large.

Counseling and Testing Data

Counseling, Testing, and Referral (CTR) data are collected and used (1) to assess the behavioral risks for sex and needle-sharing partners of HIV-diagnosed persons; (2) to evaluate the effectiveness of the CTR program as part of the overall HIV prevention effort; and (3) to improve how other HIV prevention activities, interventions, and services are implemented.

Accurate and consistent data collection is a critical component for evaluating how effective the CTR program is, as well as enabling providers to better focus prevention efforts on those persons most at risk. The data reveal information of the dynamics of HIV transmission in general, and it allows for more intensive prevention and education efforts to be applied for specific high-risk groups. To do all this, however, the collected data must be relevant to behavioral risks, HIV/AIDS prevalence and the demographics of affected communities.

The following numbers represent all tests that were administered from CTR services during the year 2007.

By the end of 2007, a total of 31,553 tests had been administered. Of those tests, 222 had positive results, which equates to a positive rate of 7.0 per 1,000 tested persons.

The group of tested persons included more males (54.9%) than females (45.0%). Table 38 lists the number of tests and the results, as well as the corresponding rates by sex.

Table 38: Number, Percentage and Rate of CTR Tests Performed by Sex in Indiana in 2007

Sex	Number of Tests	Number of Positives	New Positives	Positive Rate /1,000
Male	17,334 (54.9%)	154 (69.4%)	101 (0.5%)	8.9
Female	14,189 (45.0%)	63 (28.4%)	37 (0.2%)	4.4
Transgendered-FTM	8 (0.0%)	0 (0.0%)	0 (0.0%)	0
Transgendered-MTF	22 (0.1%)	5 (2.2%)	3 (13.6%)	227.3
Total	31,553	222	141	7.0

The rate for males and females is calculated by dividing the number of male and female positives by their respective total tested population times 1,000.

Consistent with the results from the Surveillance Report, males have a higher number of positive results than females. Still, the male positivity rate is about two times the female rate. New Positives represent those that tested that were not repeat testers. Persons often repeat test because of loss to follow-up or denial in accepting the disease.

The racial and ethnic distribution of the CTR results is shown in Table 39 for 2007.

Table 39: Number, Percentage and Rate of CTR Tests Performed by Race/Ethnicity in Indiana, 2007

Category	Number of Tests	Positives	New Positives	Positive Rate/1,000
White	17,469 (55.4%)	94 (42.3%)	64	5.4
Black	11,852 (37.5%)	117 (52.7%)	71	9.9
Other	2,232 (7.1%)	11 (5.0%)	6	4.9
Total	31,553 (100.0%)	222 (100%)	141	7.0
Hispanic*	3,072 (9.7%)	15 (6.8%)	7	4.9

* Please note that persons of Hispanic ethnicity can be of any race and are therefore not treated as a race in the Table 40.

Among the tested population, Blacks have the highest positivity rate of 9.9 per 1,000 tested persons, followed by Whites with 5.4 per 1,000. Whites make up more than half of the tested persons but less than half of all positive test results. In the last Census of the general population about 4% of Indiana's population identified themselves as Hispanic/Latino. In the group of tested persons 6.8% identified as Hispanic.

Age wise, the majority of positive results were found in the 30 to 39 year old group (11.6/1,000 tested persons). Figure 49 shows the positive rates by all age groups.

Figure 49: CTR New Positivity Rates per 1,000 by Age Group in Indiana, 2007




Table 40 lists the number of tested persons, the number of positive results and the positivity rate by age group.

Table 40: Number, Percentage and Rate of CTR Tests Performed by Age Group in Indiana, 2007

Age Group in Years	Number of Tests	Percent	Number of Positive Results		New Positives	Positive Rate/1,000
			Results	Percent		
0-12	122	0.4%	0	0	0	0
13-19	4,027	12.8%	14	6.3%	9	3.5
20-29	13,546	42.9%	74	33.3%	46	5.5
30-39	6,794	21.5%	79	35.6%	46	11.6
40-49	4,607	14.6%	44	19.8%	31	9.6
50+	2,457	7.8%	11	5.0%	9	4.5
Total	31,553	100.0%	222	100.0%	141	7.0

The majority of tests were performed among the group of 20 to 29 year olds (42.9%). The largest number of positive results occurred among the 30-39 age group 35.6% of all positive results. The 30-39 age group also had the highest positivity rate of all age groups with 11.6 per 1,000 tested persons.

An extended set of risk category variables was collected for each person tested. The results are shown in Table 41.

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007

	Incarcerated	Percent	Sex Worker	Percent
Yes	2,909	9.2%	364	1.2%
No	28,539	90.5%	31,061	98.4%
Don't Know	36	0.1%	34	0.1%
Not Asked	0	0.0%	0	0.0%
No Response	69	0.2%	94	0.3%
Blank	0	0.0%	0	0.0%
Total	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

	IDU	Percent	Sex w/ Female	Percent	Sex w/ hemophilia	Percent	Sex while on Drugs	Percent
Within 30 Days	229	0.7%	11,169	35.4%	68	0.2%	3,729	11.8%
Within 60 Days	97	0.3%	1,020	3.2%	17	0.1%	793	2.5%
Within 90 Days	73	0.2%	799	2.5%	8	0.0%	603	1.9%
Don't Know	44	0.1%	67	0.2%	810	2.6%	87	0.3%
More than 90 Days	923	2.9%	3,267	10.4%	140	0.4%	3,544	11.2%
No	30,170	95.6%	15,203	48.2%	30,477	96.6%	22,774	72.2%
No Response	17	0.1%	28	0.1%	33	0.1%	23	0.1%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

	Sex while on Heroin	Percent	Sex while on Crack	Percent	Sex while on Cocaine	Percent
Within 30 Days	184	0.6%	777	2.5%	699	2.2%
Within 60 Days	68	0.2%	235	0.7%	253	0.8%
Within 90 Days	52	0.2%	177	0.6%	195	0.6%
Don't Know	46	0.2%	56	0.2%	70	0.2%
More than 90 Days	614	2.0%	1,338	4.2%	1,962	6.2%
No	30,566	96.9%	28,947	91.7%	28,350	89.9%
No Response	23	0.1%	23	0.1%	24	0.1%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

	Sex while on Crystal Meth	Percent	Sex while on Ecstasy	Percent	Sex while on Ketamine	Percent
Within 30 Days	160	0.5%	169	0.5%	12	0.0%
Within 60 Days	80	0.3%	74	0.2%	7	0.0%
Within 90 Days	73	0.2%	101	0.3%	12	0.0%
Don't Know More than 90 Days	41	0.1%	45	0.1%	52	0.2%
No	1,047	3.3%	919	2.9%	229	0.7%
No Response	30,126	95.5%	30,213	95.8%	31,214	98.9%
	26	0.1%	32	0.1%	27	0.1%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

	Sex w/ Transgender	Percent	Sex w/ Male	Percent	Sex for Drugs/ Money	Percent
Within 30 Days	54	0.2%	12,258	38.9%	525	1.7%
Within 60 Days	9	0.0%	1,258	4.0%	155	0.5%
Within 90 Days	15	0.1%	802	2.5%	99	0.3%
Don't Know	50	0.2%	32	0.1%	67	0.2%
More than 90 Days	89	0.3%	2,559	8.1%	700	2.2%
No	31,319	99.3%	14,607	46.3%	29,986	95.0%
No Response	17	0.1%	37	0.1%	21	0.1%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

Response	Sex w/ IDU	Percent	Sex w/ HIV unknown	Percent	Sex w/ MSM	Percent	Sex w/ met from Internet	Percent
Within 30 Days	253	0.8%	10,560	33.5%	1,797	5.7%	596	1.9%
Within 60 Days	79	0.3%	1,234	3.9%	262	0.8%	157	0.5%
Within 90 Days	72	0.2%	762	2.4%	191	0.6%	106	0.3%
Do not know	1,034	3.3%	1,922	6.1%	746	2.4%	81	0.3%
More than 90 Days	948	3.0%	3,657	11.6%	673	2.1%	741	2.4%
No	29,139	92.4%	13,384	42.4%	27,844	88.3%	29,842	94.6%
No Response	28	0.1%	34	0.1%	40	0.1%	30	0.1%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 41: Number and Percentage of CTR Tests Performed by Mode of Transmission in Indiana, 2007 - continued

	Sex w/ HIV+	Percent	Sex w/ person who exchanges sex for drugs/money	Percent	Sex w/ Anonymous Partner	Percent	Other	Percent
Within 30 Days	397	1.3%	559	1.8%	1,929	6.1%	2,932	9.3%
Within 60 Days	101	0.3%	152	0.5%	560	1.8%	327	1.0%
Within 90 Days	80	0.3%	101	0.3%	414	1.3%	225	0.7%
Do not know	3,809	12.1%	771	2.4%	193	0.6%	195	0.6%
More than 90 Days	463	1.5%	823	2.6%	3,457	11.0%	1,267	4.0%
No	26,667	84.5%	29,118	92.3%	24,966	79.1%	26,547	84.1%
No Response	36	0.1%	29	0.1%	34	0.1%	60	0.2%
Total	31,553	100.0%	31,553	100.0%	31,553	100.0%	31,553	100.0%

Table 42: Number and Percentage of Confirmed STD Status of Clients, 2007

Response	Frequency	Percent
Do not know	111	0.4%
No	29,897	94.8%
Not Asked	7	0.0%
Yes- Lab Confirmed	1,035	3.3%
Yes- Self Report	503	1.6%
Total	31,553	100.0

The majority of tested clients showed that they did not have an STD (94.8%). A total of 4.9% of the clients tested showed that they had an STD, either self-reported or laboratory confirmed.

Throughout the state of Indiana HIV tests are performed at various different sites. Table 43 lists the number of tests and the different types of sites at which they were performed.

Table 43: Number and Percentage of CTR Tests Performed by Site Type in Indiana, 2007

Test Site	Number of Tests	Percent
Community Setting	516	1.6%
Community Setting – AIDS Service Organization (Non-clinical)	1,593	5.1%
Community Setting – Church/Mosque/Synagogue/Temple	274	0.9%
Community Setting – Community Center	191	0.6%
Community Setting - Bar/Club/Adult entertainment	295	0.9%
Community Setting – Other	1,367	4.3%
Community Setting – Residential	39	0.1%
Community Setting - Shelter/Transitional housing	134	0.4%
Correctional Facility	564	1.8%
Facility Other	241	0.8%
HIV Counseling & Testing Site	1,729	5.5%
Inpatient - Drug/Alcohol Treatment	395	1.3%
Inpatient Facility	43	0.1%
No Location/Response	1,478	4.7%
Community Setting	516	1.6%
Community Setting – AIDS Service Organization (Non-clinical)	1,593	5.1%
Community Setting – Church/Mosque/Synagogue/Temple	274	0.9%
Community Setting – Community Center	191	0.6%
Outpatient - Community Health Clinic	1,756	5.6%
Outpatient - Community Mental Health	810	2.6%
Outpatient - Family Planning	1,390	4.4%
Outpatient - Health Department/Public Health Clinic	16,391	51.9%
Outpatient - HIV Specialty Clinic	898	2.8%
Outpatient - School/University Clinic	165	0.5%
Outpatient – Other or Unknown	1,514	4.8%
No Label	616	2.0%
Total	31,553	100.0%

The majority of tests (51.9%) were performed at Health Departments/Public Health Clinics throughout the state.